
Editorial Board

Editorial and Advisory Board

Matsyura O.V

Altai State University, Barnaul, Russia

Deputy Chief Editor

Brygadyrenko V.V

Oles Honchar Dnipropetrovsk National University, Dnipropetrovsk, Ukraine

Yakovlev R.V

Altai State University, Barnaul, Russia

Shmakov A.I

Altai State University, Head South-Siberian Botanical Garden, Russia

Milan Vogrin

Slovenian Bird Research and Nature Protection (DPPVN), Slovenia

Eugene Maltsev, PhD

Professor

Melitopol Bogdan Khmelnytskyi State Pedagogical University, Getmanska, Melitopol, Ukraine

Jan Van Der Winden

Bureau Vaardenburg, Netherlands

Editorial Board

Mytiai I.S, PhD

Associate Professor, Animal Biology, National University of Bioresources and Nature Management of Ukraine, Ukraine

Jizhou Jiang

Distinguished Professor, School of Environmental Ecology and Biological Engineering, Wuhan Institute of Technology
Wuhan 430205, P.R. China

Professor Dr Hab. Janusz Uchmanski

Centre for Ecological Research. Polish Academy of Sciences
Poland

Dr. Slawomir Ligeza

Institute of Soil Science Environment Engineering and Management
University of Life Sciences in Lublin, Poland

Dr. Golawski Artur

Siedlce University of Natural Sciences and Humanities, Siedlce, Poland

Dr. Seyed Mehdi Talebi

Department of Biology, Arak University, Iran

Dr. Kasprzykowski Zbigniew

Siedlce University of Natural Sciences and Humanities, Siedlce, Poland

Editorial Note for Ukrainian Journal of Ecology



1 - 1

[Sam Mathew*](#)

[HTML](#)

Securement of the optimal microclimate parameters in accomodations of the educational institutions for children by implementation of the system of local air recuperation



1 - 7

[P.H. Plieshkov](#), [K.H. Petrova](#), [I.V. Savelenko*](#), [O.I. Sirikov](#) and [N.Yu. Harasova](#)

DOI: [10.15421/2020_55](#)

[Abstract](#) [HTML](#)

Bioenergy and economic efficiency technological methods growing of switchgrass



8 - 15

[V.A. Mazur](#), [Y.Y. Branitskyi](#) and [H.V. Pantsyreva*](#)

DOI: [10.15421/2020_56](#)

[Abstract](#) [HTML](#)

The State of the Spruce Stands of the Boreal and Boreal Subboreal Forests of the Eastern European Plain in the Territory of the Udmurt Republic (Russia)



16 - 22

[I. L. Bukharina*](#), [A. Konopkova](#), [K. E. Vedernikov](#), [O. A. Svetlakova](#) and [A. S. Pashkova](#)

DOI: [10.15421/2020_57](#)

[Abstract](#) [HTML](#)

Agricultural use of peat bog soils and changes in water duty to floodplains of rivers of the Left-Bank Forest-Steppe



23 - 27

[V.F. Kaminsky*](#), [N.I. Shtakal](#), [Kolomiets](#) and [V.N. Shtakal](#)

DOI: [10.15421/2020_58](#)

[Abstract](#) [HTML](#)

Influence of mineral fertilization level on productivity of *Camelina sativa* in the conditions of Prycarpattia



28 - 32

[Ya.Ya. Hryhoriv](#), [S.O. Butenko*](#), [I.M. Masyk](#), [V.I. Onychko](#), [T.O. Onychko](#), [O.I. Pshychenko](#), [V.I. Komar](#) and [O.P. Berezniak](#)

DOI: [10.15421/2020_59](#)

[Abstract](#) [HTML](#)

Assessments of physico-chemical parameters of Garaet Hadj Tahar wetland and their effect on waterbirds settlement



33 - 39

[Y. Bara](#), [M. Bara*](#), [M. Bensouilah](#), [M. Saheb](#), [S. Atoussi](#) and [M. Houhamdi](#)

DOI: [10.15421/2020_60](#)

[Abstract](#) [HTML](#)

Influence of foliar feeding on the grain productivity of corn hybrids in the conditions of the right-bank forest-steppe of Ukraine



[O.M. Kolisnyk](#), [O.O. Khodanitska](#), [A.O. Butenko*](#), [N.A. Lebedieva](#), [L.A. Yakovets](#), [O.M. Tkachenko](#), [O.L. Ihnatieva](#) and [O.V. Kurinnyi](#)

DOI: 10.15421/2020_61

[Abstract](#) [HTML](#)

Ecological and geographical characteristics of aujeszky's disease among the wild pigs in Ukraine



[V.V. Ukhovskiy*](#), [Z.M. Drozhzhe](#), [M.V. Bezymennyi](#), [N.A. Mezhenka](#), [O.M. Romanov](#), [M.S. Karpulenko](#), [O.M. Zhovnrir](#) and [A.O. Mezhenkiy](#)

[Abstract](#)

Efficiency of potato and garlic virus control by interferon use



[O. Melnyk*](#), [N. Chefonova](#), [Y. Zelendin](#), [S. Datsenko](#), [S. Shcherbina](#), [O. Vitanov](#), [O.Semenchenko](#), [N. Dukhina](#), [I. Mytenko](#) and [D. Ivanin](#)

[Abstract](#)

Ecological plasticity and productive potential of tobacco in Central Forest Steppe of Ukraine



[K.P. Leonova*](#), [A.V. Morgun](#), [H.M. Hospodarenko](#), [I.Y. Rassadina](#), [V.G. Kryzhanovskiy](#) and [A.T. Martyniuk](#)

DOI: 10.15421/2020_64

[Abstract](#) [HTML](#)

Comparative histochemical analysis of phosphatases activity of the skin glands in some artiodactyls



[A.B. Kiladze*](#) and [N.K. Dzhemukhadze](#)

DOI: 10.15421/2020_65

[Abstract](#) [HTML](#)

A Study of the growth and yield of *Solanum lycopersicum* under greenhouses differentiated by the LDPE cover-film



[A. Belhachemi*](#), [M. Maatoug](#), [M. Amirat](#) and [A. Dehbi](#)

DOI: 10.15421/2020_66

[Abstract](#) [HTML](#)

Effect of antibiotics, hormones and anthelmintic on high molecular weight protein fractions in the common carp



[I. Kurbatova*](#), [M. Zakharenko](#), [O. Tupitska](#), [O. Yaremchuk](#) and [L. Chepil](#)

DOI: 10.15421/2020_67

[Abstract](#) [HTML](#)

Growth and productivity of winter wheat (*Triticum aestivum* L.) depending on the sowing parameters




[S. Poltoretskyi*](#), [S. Tretiakova](#), [I. Mostoviak](#), [A. Yatsenko](#), [Y. Tereshchenko](#), [N. Poltoretska](#) and [A. Berezovskyi](#)

DOI: 10.15421/2020_68

[Abstract](#) [HTML](#)

Taxonomic study of family Asteraceae from ShishiKoh Valley,
Chitral, Pakistan


 88 - 91

 [K. Ullah*](#), [G.M. Shah](#), [J. Alam](#) and [M. Hussain](#)

 Abstract


Ethnobotany of the Medicinal Plants Used by Indigenous
Communities in the Mountain of Shishikoh Valley, Hindukush Chitral

 92 - 105

 [K. Ullah*](#), [G.M. Shah](#), [J. Alam](#) and [M. Hussain](#)

 Abstract

Effect of 6-Benzylaminopurine on Morphogenesis and Production
Process of Sweet Pepper (*Capsicum annum L.*)

 106 - 111


 [V.G. Kuryata*](#), [O.V. Kushnir](#) and [O.O. Kravets](#)

 Abstract


Species Composition and Noxiousness of Segetal Vegetation in
Winter Rye Agrocoenoses in the Central Ukrainian Polissia


 112 - 117


 [M.M. Kliuchevych](#), [S.H. Stoliar*](#), [O.Yu. Hrytsenko](#), [S.V. Retman](#), [H.M. Tkalenko](#) and [L.V. Bilotserkivska](#)

 Abstract


The Study of The Level of Environmental Consciousness and the
Efficiency of Environmental Education Students

 118 - 125

 [I. Ivanov*](#), [L. Frolova](#), [N. Prokopenko](#), [O. Belyanovskaya](#) and [O. Prokopenko](#)

 Abstract


Changes in the biochemical composition of the seed material of
sunflower hybrids during long-term storage


 126 - 130

 [O.A. Yeremenko*](#), [L.?. Pokoptseva](#), [L.V. ??dorova](#) and [?.V. Shepel](#)

 Abstract


Intensity of heavy metal accumulation in plants of *Silybum*
marianum L. in conditions of field rotation


 131 - 136


 [S.F. Razanov](#), [O.P. Tkachuk*](#), [A.M. Razanova](#), [M.I. Bakhmat](#) and [O.M. Bakhmat](#)

 Abstract

The activity of the microbial groups of maize root-zone in
different crop rotations

 137 - 140

 [O.Yu. Karpenko](#), [V.M. Rozhko](#), [A.I. Lychuk](#), [A.O. Butenko*](#), [G.A. Davydenko](#), [D.S. Tymchuk](#), [O.L. Tonkha](#) and [V.P. Kovalenko](#)

 Abstract

Allelopathic Activity of Secretions of Plant Mass and Soil From the
Form of Monard Double (*Monarda didyma L.*)


 141 - 145


 [N.V. Zaimenko*](#), [N.O. Hnatyuk](#) and [B.O. Ivanytska](#)

 Abstract

Empowering informal settlements around metropolises (Case
study: District 19 of Tehran, Iran)


 156 - 164

 [S. Zeynab*](#)

 Abstract


Specific activity of Sr-90 and Cs-137 in rabbits of various genotypes


 165 - 169

 O.V. Bojko, L.M. Darmohray, I.S. Luchyn, O.F. Honchar and B.V. Gutyj*

 Abstract


Analysis of spatial and temporal dynamics of epizootic process of blackleg in Ukraine


 170 - 176

 P.K. Boiko*, V.M. Sokolyuk, O.P. Boiko, V.I. Koziy and A.M. Fedorchenko

 Abstract

Phylogeny of the North Asian Cystopteridaceae (Polypodiopsida) based on trnG-R intergenic spacer


 177 - 183

 I.I. Gureyeva*, D.O. Ulko*, R.S. Romanets and A.A. Kuznetsov

 Abstract


Effect of reservoir temperature and oxygen conditions on the activity of Na-K pump in embryos and larvae of perch, roach, and ruffe


 184 - 189

 O. Vodianskyi, O. Dadak, V. Maxym, V. Dushka, B. Gutyj*, L. Zagoruy, A. Sliusarenko, O. Rud, N. Prysiazniuk, Y. Khudiyash, O. Khomiak, N. Hrynevych, O. Potrokhov and V. Liublin

 Abstract


Experience of developed countries in environmental safety policy

 190 - 194

 A. I. Krukov, O. V. Radchenko, O. O. Radchenko, B.K. Garmash, Ye.S. Biletska, R.V. Ponomarenko, S. I. Sysoieva, S. V. Stankevych* and S.O. Vynohradenko

 Abstract


Long-term monitoring of arable lands infestation in the steppe zone of Ukraine


 195 - 199


 O.N. Kurdyukova*

 Abstract


A Review of germanium environmental distribution, migration and accumulation

 200 - 208


 O.I. Sobolev, A.O. Vysotskij, N.D. Levkivska, H.V. Sus, V.B. Todoriuk, M.M. Zhelavskiy, O.Y. Petryshak, V.I. Kushnir, O.S. Naumyuk, R.A. Petryshak, I. M. Kushnir, O.O. Borshch, S.V. Soboliev?, B.V. Gutyj* and N.V. Magrelo

 Abstract


A Study of the Emerald Network objects in Ukrainian Forest-Steppe of Dnieper Ecological Corridor


 209 - 218

 I.V. Solomakha, V.V. Konishchuk, O.V. Mudrak* and H.V. Mudrak

 Abstract

Morphobioecological features and harmfulness of apple-blossom weevil (*Anthonomus pomorum* Linnaeus, 1758)

 219 - 230

 I.V. Zabrodina*, T.A. Romanova, O.V. Romanov, V.O. Melenti, D.D. Yushchuk, L.Ya. Sirous, M.O. Filatov, I.P. Lezhenina, H.V. Baidyk, O.A. Molchanova, S.V. Stankevych, M.D. Yevtushenko and O.M. Bragin

 Abstract


Effects of selenium compounds and toxicant action on oxidative biomarkers in quails



232 - 239


 S.I. Tsekhmistrenko*, O.P. Shulko, H.V. Merzlova, Yu.O. Melnichenko, M.M. Fedorchenko, N.V. Rol, S.A. Polishchuk, N.V. Ponomarenko, N.O. Tymoshok, V.M. Kharchyshyn, O.M. Melnichenko, O.S. Tsekhmistrenko, V.S. Bityutskyy and A.A. Demchenko

 Abstract


Probability of Radon cancer occurrence in Rivne city residents (Ukraine), calculation by WISMUT and Radon-2014 models



240 - 248


 O.O. Lebed*, L.V. Klymenko, V.S. Trotsyuk, V.A. Hulchuk and A.V. Lysytsya

 Abstract


Carbon Absorption Ability of Pine Forest Plantations in the Ukrainian Polissya


249 - 255


 V.V. Moroz*, Y.A. Nykytiuk, P.A. Nykytiuk, M.M. Kliuchevych and O.M. Komorna

 Abstract


The Immunological Reactivity of Turkeys of Different Genotypes on the Action of Environmental Factors



256 - 259

 Ya. Rebezov, F. Khaziakhmetov, N. Fedoseeva, I. Knysh, N. Vinogradova, S. Safronov, M. Derkho, T. Bezhinari, M. Rebezov*, O. Gorelik and A. Khabirov

 Abstract


Influence of Transport Stress on the Adaptation Potential of Chickens



260 - 263

 O. Gorelik, M. Ali Shariati, T. Bezhinari, N. Lop?eva, P. Burkov, S. Ermolov, I. Knysh, N. Vinogradova, M. Eliseenkova, I. Dolmatova, M. Derkho, S. Harlap and M. Rebezov*

 Abstract


Dynamics of Hematological Indicators of Chickens under Stress-Inducing Influence


264 - 267


 O. Gorelik, S. Harlap, N. Lop?eva, T. Bezhinari, V. Kosilov, P. Burkov, I. Ivanova, S. Gritsenko, I. Dolmatova, O. Tsareva, S. Safronov, M. Ali Shariati and M. Rebezov*

 Abstract

Productivity of Sugar Maize of Hybrid Moreland F1 Depending on Technological Factors of Growing



268 - 272

 Ya.Ya. Hryhoriv, A.O. Butenko*, G.A. Davydenko, M.V. Radchenko, O.M. Tykhonova, L.V. Kriuchko and Z.I. Hlupak

 Abstract


Climate Change: Apocalyptic Prognosis and Reality


273 - 278

 V.S. Bondar, A.V. Fursa*, M.Ya. Gumentyk and I.V. Svystunova

 Abstract

Growth regulatory Activity of 2-Methyl-4-Thioquinoline Derivatives


279 - 283

 M.M. Komet, G.F. Dudareva, V.I. Gencheva*, O.O. Klimova, V.V. Peretiatko and O.A. Brazhko

 Abstract

Economic and Energy Efficiency of Forming and Using Legume-Cereal Grass Stands Depending on Fertilizers



284 - 288

[U.M. Karbivska](#), [V.G. Kurgak](#), [V.F. Kaminskyj](#), [A.O. Butenko*](#), [G.A. Davydenko](#), [O.B. Viunenko](#), [S.M. Vyhanialo](#) and [S.V. Khomenko](#)

Abstract

Efficiency of premixes with Bioplex microelements in the diets of Holstein cattle



289 - 293

[A. Horchanok*](#), [O. Kuzmenko](#), [L. Lytvshchenko](#), [M. Lieshchova](#), [N. Prysiazhniuk](#), [O. Bezv](#) and [O. Slobodeniuk](#)

Abstract

Treatment strategies for sheep with acute yellow atrophy of the liver caused by the fasciolosis



294 - 301

[B.O. Chemushkin*](#), [H.O. Zinko](#), [B.O. Lukashchuk](#), [V.L. Fedorovych](#), [M.H. Lychuk](#), [V.I. Rusyn](#), [M.I. Leno](#), [I.A. Maksymovych](#), [A.R. Shcherbatyy](#), [B.V. Gutyj](#), [L.G. Slivinska](#), [V.V. Viizlo](#) and [O.I. Prystupa](#)

Abstract

Effect of *Echinacea pallida* supplementation on the amino acid and fatty acid composition of Pharaoh Quail meat



302 - 307

[R.A. Chudak*](#), [V.M. Ush?kov](#), [Y.M. Poberezhets](#), [H.I. Lotka](#), [?V. Polishchuk](#) and [L.V. Kazmiruk](#)

Abstract

Radon induced lung cancer in residents of Rivne (Ukraine) calculated by various risk models



311 - 318

[O.O. Lebed*](#), [A.V. Lysytsya](#), [V.O. Myslinchuk](#), [L.V. Hladun](#) and [A.V. Rybalko](#)

Abstract

Annotated key to weevils of the world. Part 4. Subfamilies *Eirrhinae*, *Dryophthorinae* and *Cossoninae* (Curculionidae)



319 - 331

[A.A. Legalov*](#)

Abstract

Annotated key to weevils of the world: Part 5 - Subfamily Entiminae (Curculionidae)



332 - 346

[A.A. Legalov*](#)

Abstract

Effect of Butaselmavit-Plus on the immune system of piglets during and after weaning



347 - 352

[T. V. Martyshuk](#), [B. V. Gutyj*](#), [M. M. Zhelavskiy](#), [S. V. Midyk](#), [A. M. Fedorchenko](#), [V. B. Todoruk](#), [T. B. Nahirniak](#), [Ya. V. Kiser](#), [H. V. Sus](#), [V. A. Chemerys](#), [N. D. Levkivska](#) and [I. I. Iglitskej](#)

Abstract

A new species of the genus *Basiliogeus* Kuschel, 1994 (Insecta: Coleoptera: Nemonychidae) from Malaysia





353 - 356


[A.A. Legalov*](#)

Abstract


Two new species and new findings of Curculionoidea (Insecta: Coleoptera) in Baltic Amber


357 - 364

 [A.A. Legalov](#) and [G.O. Poinar, Jr.*](#)

 [Abstract](#)


Development and reproductive qualities of sows of different breeds: innovative and traditional methods of assessment



365 - 370

 [V. Khalak*](#), [B. Gutyj](#), [O. Bordun](#), [A. Horchanok](#), [M. Ilchenko](#), [S. Smyslov](#), [O. Kuzmenko](#) and [L. Lytyshchenko](#)

 [Abstract](#)

Effect of morphological and biochemical blood composition on milk yield in Simmental breed cows of different production types



371 - 377

 [N.P. Mazur](#), [V.Yu. Hudyma](#), [N.P. Stakhiv](#), [H.H. Denys](#), [O.S. Grabovska](#), [T.V. Orikhivskiy](#), [N.M. Kuziv](#), [M.I. Kuziv](#), [B.V. Gutyj*](#), [P.V. Bodnar](#), [O.V. Fedorovych](#), [E.I. Fedorovych](#), [V.V. Fedorovych](#) and [N.I. Pakholkiv](#)

 [Abstract](#)

Improving the sanitary condition of fish pond bed by forage grass cultivation



378 - 384


 [S.M. Nazarenko](#), [A.P. Paliy](#), [J.V. Negreba](#), [L.N. Lazorenko](#), [O.I. Kasianenko](#), [R.V. Petrov](#), [O.V. Fotin](#), [A.I. Fotin](#), [A.V. Berezovskiy](#), [A.P. Paliy*](#) and [H.I. Rebenko](#)

 [Abstract](#)

Milking and udder health assessment in industrial farming



385 - 391

 [A.P. Paliy*](#), [D.A. Levkin](#), [V.B. Savchenko](#), [A.I. Sychov](#), [T.Y. Mitiashkina](#), [S.A. Denicenکو](#), [O.V. Bogomolov](#), [V.V. Bredykhin](#), [Y.E. Rozum](#), [A.P. Reshetnichenko](#), [H.F. Chechui](#), [S.A. Mihalchenko](#) and [A.P. Paliy](#)

 [Abstract](#)


Microbial contamination of cow's milk and operator hygiene



392 - 397

 [A.P. Paliy*](#), [O.M. Pobirchenko](#), [O.V. Kovalenko](#), [V.S. Kalabska](#), [Y.O. Kovalchuk](#), [I.S. Pavlenko](#), [M.V. Kazakov](#), [V.M. Borovkova](#), [L.L. Kushch](#), [S.A. Zolotaryova](#), [K.O. Rodionova](#), [A.P. Paliy](#) and [O.S. Umrihina](#)

 [Abstract](#)


Improved techniques for the reduction of microbial contamination toxic gas content in the air of the poultry houses



398 - 403

 [A.P. Paliy*](#), [Y.A. Boyko](#), [A.O. Kovalchuk](#), [I.V. Grebnova](#), [Yu.I. Tokolov](#), [A.M. Milenin](#), [I.V. Chaly](#), [O.A. Svirgun](#), [P.V. Gurskiy](#), [I.M. Lukyanov](#) and [A.P. Paliy](#)

 [Abstract](#)

Microstructural analysis of sausage quality


404 - 409

 [?P. Paliy*](#), [T.M. Ihnatieva](#), [T.V. Dudus](#), [H.V. Ovcharenko](#), [N.I. Sakhniuk](#), [Ye.V. Vashchuk](#), [N.M. Bogatko](#), [K.O. Rodionova](#), [A.P. Paliy](#), [B.T. Stegnyy](#) and [L.V. Kovalenko](#)

 [Abstract](#)

The estimation of metagenome and functionally polymorphisms of soil prokaryote



410 - 414

[O.T. Piven*](#), [M.S. Khimych](#), [V.Z. Salata](#), [B.V. Gutyj](#), [O.V. Naidich](#), [H.A. Skrypka](#), [Z.B. Koreneva](#), [I.V. Dvylyuk](#), [O.M. Gorobey](#) and [V.O. Rud](#)

Abstract

Contmination of heavy metals and radionuclides in the honey with different production origin



415 - 419

[O.T. Piven*](#), [O.V. Ivleva](#), [L.I. Nalyvayko](#), [V.F. Mohutova](#), [N.M. Bogatko](#), [T.I. Fotina](#), [I.V. Yatsenko](#), [V.Z. Salata](#), [M.S. Khimych](#) and [T.A. Odyntsova](#)

Abstract

Effect of ultraviolet irradiation on beef carcass yield



420 - 425

[L. Roman](#), [M. Broshkov](#), [I. Popova](#), [A. Hierdieva](#), [S. Sidashova](#), [N. Bogach](#), [S. Ulizko](#) and [B. Gutyj*](#)

Abstract

Influence of ovarian follicular cysts on reproductive performance in the cattle of new Ukrainian red dairy breed



426 - 434

[I. Simonova](#), [S. Grabovskyi*](#), [U. Drachuk](#), [B. Halukh](#) and [I. Basarab](#)

Abstract

Amino acid composition of meat and bone meal from various manufacturers of pet food and animal feed



435 - 439

[I. Simonova](#), [S. Grabovskyi*](#), [U. Drachuk](#), [B. Halukh](#) and [I. Basarab](#)

Abstract

Separate indices of homeostasis and the balance of the prooxidant-oxidant system in sheep for fetoplacental insufficiency



440 - 448

[P.M. Skliarov*](#), [S.J. Fedorenko](#), [S.V. Naumenko](#), [O.V. Onyshhenko](#), [A.M. Pasternak](#) and [K.O. Holda](#)

Abstract

Evaluation of the tobacco genotypes by seed productivity



449 - 454

[K. P. Leonova*](#), [A. V. Morgun](#), [H. M. Hospodarenko](#), [V. V. Ketskalo](#), [S. P. Kotsyuba](#) and [V. I. Nevlad](#)

Abstract

Change of water properties during the process of floodplain vegetation destruction



455 - 460

[M.S. Bobrova](#), [I.V. Luschaeva](#), [L.F. Shepeleva](#), [O.S. Pokrovsky](#), [E.S. Rabtsevich](#), [L.P. Borilo](#), [E. Lutova](#), [A.S. Prokushkin](#), [V.Yu. Shuvarikova](#), [R.M. Manasypov](#) and [L.G. Kolesnichenko*](#)

Abstract

Effect of *Echinacea pallida* supplementation on the amino acid and fatty acid composition of Pharaoh Quail meat

R.A. Chudak*, V.M. Ushakov, Y.M. Poberezhets, H.I. Lotka, T.V. Polishchuk, L.V. Kazmiruk

Vinnitsia National Agrarian University, 3 Soniachna St., Vinnitsia, 21008, Ukraine

*Corresponding author E-mail: romanchudak@ukr.net

Received: 13.04.2020. Accepted 23.05.2020

The purpose of the research was to investigate the effect of *Echinacea pallida* extract effect on the amino acid and fatty acid content of Pharaoh Quail meat. We proved that the amount of essential amino acids was greater by 1.31% in the pectoral muscles of Quails that consumed 12 mg/kg of *Echinacea pallida* dry extract. This increase was caused by the better accumulation of lysine, arginine, threonine, leucine, and phenylalanine. The amount of essential amino acids was the highest in the thigh muscles of the control group. We found that the quality of fatty acid composition of the pectoral muscles was better due to the greater accumulation of linoleic, γ -linolenic, and α -linolenic acids. The accumulation rate of linoleic, γ -linolenic, and arachidonic fatty acids was higher in thigh muscles of the experimental Quail.

Keywords: Quails; Phytobiotics; *Echinacea pallida*; Amino acids; Meat; Fat acids

Introduction

In recent years, feed additives of natural origin have been extensively used in animal husbandry. Their wide usage explained by the accessibility, absence of adverse reactions, and the wide range of positive biological effects (Cross & Hillman, 2004; Grashorn, 2010; Podolian, 2016.). The antibiotics' application in European countries has been forbidden. This triggered the search for natural additives of natural origin containing biologically active substances that increase productivity, strengthen immunity and improve digestive processes (Ibatullin *et al.*, 2017; Podolian, 2017).

Phytobiotics are increasingly used in animal feeding. They are the phytocorrectors modifying the digestive glands work and providing the conditions for competitive growth of beneficial microflora, which stabilizes acidity and enhances the processes of nutrient absorption (Bauer & Wagner, 1991; Chudak *et al.*, 2018; Magomedaliev *et al.*, 2019). Feed additives of plant origin included into the farm animals diets help to increase productivity; improve the gut microflora; improve the taste of feed; stimulate salivation and secretion of digestive juices due to the rapid passage of feed and absorption of nutrients, and the nutrient system (Aituan & Wanyu, 2009; Bohmer & Salisch 2009).

Recently, foreign and domestic scientists have become interested in the study and application of the genus *Echinacea* species (Bauer *et al.*, 1988; Dehkordi & Fallah, 2011). They have very valuable medicinal, fodder properties, and their chemical composition contains a large number of biologically active substances. They are carbohydrates, nitrogenous substances, organic acids, lipids, vitamins, polyphenolic compounds (coffee derivatives, flavonoids, and tannins), isoprenoids, saponins, essential oils, alkaloids and minerals (Nazir & Grashorn, 2010; Barnes *et al.*, 2005; Samorodov *et al.*, 2013).

Echinacea pallida is becoming very popular in veterinary as therapeutic and prophylactic medicine and plant feed additive (Allen, 2003; Diakonova 2009; Chudak *et al.*, 2019). It has a mild action, low toxicity, and high content of vital substances affecting the animals' metabolism (Bauer & Wagner, 1990; Mamchur *et al.*, 1993; Pospelov *et al.*, 2007).

The purpose of the research was to investigate the effect of *Echinacea pallida* extract on the fatty acid and amino acid content of the Pharaoh Quail meat.

Methods

We performed the study of *Echinacea pallida* feed additive at the research farm of Vinnitsia National Agrarian University. The scientific experiment was conducted on poultry by groups-analogues method taking into account live weight, age, sex, breed, productivity, and conditions of keeping and feeding (Ibatullin *et al.*, 2017). The duration of the experiment was 56 days. Four groups of one-day-old Pharaoh meat Quail were selected for the experiment, 50 birds in each group. The control group consumed a basic diet (BD), i.e., complete feeds of the Multigain trademark. The experimental groups were additionally fed by different doses of *Echinacea pallida* feed additive extract (Table 1).

Table 1. Feeding patterns.

Group	Feeding characteristics
Control	BD (complete feeds)
II	BD + <i>Echinacea pallida</i> (6 mg/kg)
III	BD + <i>Echinacea pallida</i> (12 mg/kg)
IV	BD + <i>Echinacea pallida</i> (18 mg/kg)

The researched phytobiotics was homogeneous, powdered and dry extract of *Echinacea pallida* roots. This was brown powder with a characteristic smell and a specific bitter taste. The manufacturer of the extract from *Echinacea pallida* is LLC Kharkiv Research Plant, Ukraine. The main active biologically substances are polysaccharides, i.e., fructosans, phenolic compounds, hydroxycoric acids having anti-inflammatory, antimicrobial and adaptogenic activities. We have researched the fatty acid composition by gas chromatography. The meat amino acid composition was determined in the Laboratory of the Biochemistry Research Institute named after O.V. Paladin (Kyiv) by TTT 339 automatic analyzer using LG ANB cation exchange resin with SO₃ active group. The Tables data are presented as mean and standard deviation, the amino acid content is calculated in percent per 100 mg.

Results and Discussion

We found that different doses of *Echinacea pallida* feed additive extract could determine the amino acid content in the Quail meat (Table 2). We also revealed that the poultry consuming the researched supplement had a higher lysine content in the white meat compared to the control. This content was by 0.45% ($P < 0.001$) higher in group III and by 0.12% ($P < 0.001$) in group IV. However, the content of amino acids in the meat of Quails from group II was lower by 0.31% ($P < 0.001$).

The highest content of histidine was in the pectoral muscles of the group II, it was higher by 0.29% ($P < 0.001$); the content of arginine in the pectoral muscles of the group III was higher by 0.37% ($P < 0.001$) compared to control data. The content of aspartic acid increased by 0.65% ($P < 0.001$) in the pectoral muscles of group III, and glutamic acid increased by 0.27% ($P < 0.001$) in the pectoral muscles of group IV.

Table 2. Amino acid composition of Quail pectoral muscles.

Amino acid	Groups			
	Control	II	III	IV
Lysine	9.29 ± 0.007	8.98 ± 0.012***	9.74 ± 0.005***	9.41 ± 0.014***
Histidine	3.26 ± 0.024	3.55 ± 0.013***	3.37 ± 0.014**	3.50 ± 0.002***
Arginine	6.64 ± 0.037	6.94 ± 0.022***	7.01 ± 0.012***	6.72 ± 0.029
Aspartic acid	6.50 ± 0.005	6.17 ± 0.003***	7.15 ± 0.012***	6.61 ± 0.010***
Threonine	4.89 ± 0.019	5.12 ± 0.014***	4.91 ± 0.009	4.48 ± .014***
Serine	4.19 ± 0.014	4.40 ± 0.003***	4.21 ± 0.009	4.20 ± 0.002
Glutamic acid	16.77±0.030	16.42±0.040***	15.38 ±0.017***	17.04±0.027***
Proline	5.06 ± 0.058	4.35 ± 0.077***	3.55 ± 0.036***	4.36 ± 0.051***
Glycine	4.72 ± 0.012	4.85 ± 0.002***	4.96 ± 0.003***	4.74 ± 0.007
Alanine	6.26 ± 0.007	6.39 ± 0.010***	6.49 ± 0.003***	6.07 ± 0.009***
Cystine	1.19 ± 0.029	1.31 ± 0.017*	1.25 ± 0.012	1.39 ± 0.015***
Valine	5.36 ± 0.025	5.56 ± 0.020***	5.60 ± 0.007***	5.52 ± 0.008***
Methionine	2.88 ± 0.011	2.75 ± 0.013***	2.89 ± 0.010	2.98 ± 0.017**
Isoleucine	5.01 ± 0.016	5.21 ± 0.030**	5.17 ± 0.014***	5.11 ± 0.007**
Leucine	9.12 ± 0.031	9.27 ± 0.035*	9.49 ± 0.007***	9.06 ± 0.020
Tyrosine	4.15 ± 0.036	4.05 ± 0.042	4.04 ± 0.012*	4.09 ± 0.030
Phenylalanine	4.65 ± 0.025	4.61 ± 0.021	4.73 ± 0.009*	4.67 ± 0.009
Essential acids amount	51.10	51.99	52.41	51.45
Substitutable acids amount	48.84	47.94	47.03	48.50

The threonine and serine content in the pectoral muscles of the Quail exceeded by 0.23% and 0.21% ($P < 0.001$) in II group fed by researched additive. It was found that the highest content of glycine and alanine was observed in II and III groups; it was higher by 0.13%, 0.24% and 0.13%, 0.23% ($P < 0.001$) respectively than control one. The cystine content increased in pectoral muscles of II and IV groups by 0.12% ($P < 0.05$) and 0.20% ($P < 0.001$) respectively than control one. It was researched that additional feeding by *Echinacea pallida* dry extract to the Quail causes increase of methionine in group IV by 0.10% ($P < 0.01$). The highest proportion

of phenylalanine was recorded in the group III; it was by 0.08% ($P < 0.05$) higher than control one. The researched additive causes lysine increasing in Quail red meat in group II by 0.68% ($P < 0.001$) respectively than control one (Table 3).

It should be mentioned that the histidine level in the third poultry group was less than the control analogues by 0.1% ($P < 0.001$). It was found that the amount of histidine and arginine decreases in the femur muscles of the experimental groups compared to the control one. The content of aspartic acid in the poultry thighs outweighed the control sample in group IV by 0.83% ($P < 0.001$) and glutamic acid by 2.26% ($P < 0.001$) in group III. The amount of threonine in the poultry muscles of the group IV increased by 0.05% ($P < 0.01$).

Table 3. Amino acid composition of Quail femur muscles.

Amino acid	Groups			
	Control	II	III	IV
Lysine	9.07 ± 0.011	9.75 ± 0.005***	9.22 ± 0.019***	9.66 ± 0.008***
Histidine	2.79 ± 0.004	2.76 ± 0.014	2.69 ± 0.008***	2.79 ± 0.007
Arginine	6.88 ± 0.004	6.57 ± 0.024***	6.71 ± 0.030**	6.73 ± 0.014***
Aspartic acid	6.60 ± 0.009	7.22 ± 0.011***	7.30 ± 0.002***	7.43 ± 0.009***
Threonine	5.08 ± 0.008	5.00 ± 0.007***	4.98 ± 0.002***	5.13 ± 0.007**
Serine	4.40 ± 0.002	4.33 ± 0.005***	4.34 ± 0.003***	4.34 ± 0.007***
Glutamic	16.93 ± 0.005	17.12 ± 0.017***	19.19 ± 0.021***	17.51 ± .021***
Proline	4.76 ± 0.031	4.86 ± 0.064	4.88 ± 0.088	4.81 ± 0.081
Glycine	5.06 ± 0.005	5.03 ± 0.005**	5.09 ± 0.002**	4.99 ± 0.008***
Alanine	5.94 ± 0.005	5.88 ± 0.008***	5.91 ± 0.012	5.76 ± 0.005***
Cystine	1.28 ± 0.012	1.25 ± 0.012	1.04 ± 0.012***	1.24 ± 0.012
Valine	5.20 ± 0.013	4.99 ± 0.005***	4.79 ± 0.012***	4.99 ± 0.010***
Methionine	3.13 ± 0.002	3.05 ± 0.009***	2.55 ± 0.012***	2.84 ± 0.012***
Isoleucine	4.94 ± 0.007	4.83 ± 0.010***	4.45 ± 0.002***	4.79 ± 0.016***
Leucine	9.05 ± 0.008	8.81 ± 0.020***	8.77 ± 0.020***	8.73 ± 0.016***
Tyrosine	4.17 ± 0.047	3.85 ± 0.023***	3.67 ± 0.017***	3.79 ± 0.019***
Phenylalanine	4.75 ± 0.078	4.61 ± 0.007	4.33 ± 0.009***	4.40 ± 0.007**
Essential acids amount	50.89	50.37	48.49	50.06
Substitutable acids amount	49.14	49.54	51.42	49.87

It was found that serine content significantly decreased in the femurs of groups II, III and IV by 0.07%, 0.06%, and 0.06%, respectively ($P < 0.001$) under the action of *Echinacea pallida* extract. Consumption of phytobiotic supplement causes the proline content increasing in red meat of groups II, III and IV by 0.1%, 0.12% and 0.05% respectively. However, a probable difference hasn't been established. The glycine content in the femur Quail muscles was greater by 0.03% ($P < 0.01$) in the group III than in the control sample. However, this indicator was lower by 0.03% ($P < 0.01$) and 0.07% ($P < 0.001$) in the groups II and IV. The level of alanine significantly decreased by 0.06% and 0.18% ($P < 0.001$) in the groups II and IV than the control group. It should be noted that feeding by *Echinacea pallida* extract different doses causes valine content decreasing by 0.21%, 0.41% and 0.21% ($P < 0.001$) in the femurs of the experimental poultry groups.

The lowest amount of cystine and methionine was in the red meat of the group III by 0.24% and 0.58% ($P < 0.001$), respectively, compared to the group I. It should be noted that the lowest levels of isoleucine and leucine were in the poultry muscles from the experimental groups II and IV, respectively by 0.49% and 0.32% ($P < 0.001$) compared to the control.

It was noted that the smallest tyrosine accumulation was observed in the femurs of the poultry of all experimental groups by 0.32%, 0.5% and 0.38% ($P < 0.001$), respectively, compared to the control sample. It was also found that the reduction of phenylalanine content is observed in the Quails of the groups III and IV by 0.42% ($P < 0.001$) and 0.35% ($P < 0.01$), respectively, compared to the control group analogues.

The amount of essential amino acids decreases by 0.52%, 2.4% and 0.83% in Quails of the experimental groups fed by the experimental supplement. However, the amount of substitutable amino acids increased by 0.4%, 2.28% and 0.73% in these groups, respectively, compared to the first poultry control group. Thus, mixed doses of *Echinacea pallida* dry extract have a positive effect on the amount of substitutable and essential amino acids in the Quail meat.

We determined that different doses of *Echinacea pallida* could influence the fatty acid content of the Quail pectoral muscles (Table 4).

Table 4. Fatty acid composition of Quail pectoral muscles (the average value in the sample).

Fatty acid	Groups			
	Control	II	III	IV
Myristic	0.29	0.25	0.30	0.22
Stearic	12.69	11.54	11.33	14.38
Palmitic	18.61	16.91	18.30	16.56
Arachidic	0.14	0.13	0.11	0.05
Margaric	0.12	0.17	0.21	0.15
Pentadecanoic	0.06	0.07	0.03	0.05
Linoleic	23.45	25.38	28.86	26.23
γ -linolenic	0.12	0.14	0.19	0.19
α -linolenic	1.05	1.20	1.20	1.26
Arachidonic	9.04	8.55	8.50	12.26
Oleic	25.92	27.40	24.27	20.11
Gondoinic	0.07	0.15	0.06	0.05
Pentadecileine	0.03	0.03	0.04	0.02
Palmitoleic	5.90	5.83	3.74	4.47
Margarinolein	0.03	0.06	0.05	0.05
Dihomolinoleic	0.12	0.17	0.07	0.15
Dihomo- γ -linolenic	0.13	0.08	0.08	0.31
Docosatetraenoic	0.09	0.08	0.08	0.13
Docosapentaenoic	0.09	0.09	0.08	0.14
Clupanodonic	0.27	0.38	0.20	0.41
Docosaheptaenoic	1.79	1.38	2.29	2.78

Table 5. Fatty acid composition of Quail femur muscles (the average value in the sample).

Fatty acid	Groups			
	Control	II	III	IV
Myristic	0.24	0.22	0.20	0.21
Stearic	16.85	14.27	17.11	17.43
Palmitic	15.47	17.30	15.59	14.32
Arachidic	0.06	0.10	0.15	0.10
Margaric	0.14	0.09	0.14	0.17
Pentadecanoic	0.05	0.04	0.02	0.02
Linoleic	23.19	24.94	25.67	25.86
γ -linolenic	0.12	0.18	0.13	0.15
α -linolenic	0.92	0.80	0.73	1.13
Arachidonic	9.54	9.93	11.93	11.32
Oleic	23.27	23.06	19.03	18.74
Gondoinic	0.06	0.12	0.09	0.12
Pentadecileine	0.01	0.02	0.02	0.02
Palmitoleic	5.62	4.47	2.90	3.47
Margarinolein	0.05	0.05	0.04	0.03
Dihomolinoleic	0.15	0.14	0.10	0.23
Dihomo- γ -linolenic	0.08	0.16	0.11	0.26
Docosatetraenoic	0.33	0.46	0.45	0.71
Docosapentaenoic	0.18	0.32	0.24	0.32
Clupanodonic	0.41	0.75	1.25	0.75
Docosaheptaenoic	3.24	2.58	4.08	4.64

The supplementary feeding of poultry (group IV) by the maximum dose of the researched feed additive caused the increase of stearic acid content in white poultry meat by 1.69%, margaric acid by 0.03%, linoleic acid by 2.78%, γ -linolenic acid by 0.07% and α -linolenic acid by 0.21% than the control one. However, the poultry of this group has decreased content of myristic, palmitic, arachidic and oleic fatty acids by 0.07%, 2.05%, 0.09% and 5.81%, respectively, compared with indicators of the group I. Minimal and medium doses of *Echinacea pallida* extract cause increased linoleic acid content by 1.93% and 5.41% in the pectoral muscle of the poultry compared to the control group. Poultry of the experimental groups II and III had a decrease of stearic and palmitic fatty acids by 1.15%, 1.36% and 1.7%, 0.31% respectively compared to the control analogues. The content of essential fatty acids such as dihomo- γ -linolenic and arachidonic acids increased by 0.18% and 3.22% in the fourth experimental group compared to the control indicators. The levels of arachidonic acid decreased by 0.49% and 0.54% respectively in the groups II and III, relative to the control data.

We found that the concentration of docosatetraenoic and docosapentaenoic acids in the pectoral muscles exceeded the control by 0.04% and 0.05% in the group IV fed by the maximum dose of the researched supplement. The content of clupanodonic and docosahexaenoic fatty acids increased in the bird of the group IV by 0.14% and 0.99%, respectively, compared with the control sample. Researches have proved that feeding different doses of *Echinacea pallida* extract changed the fatty acid composition of the femur muscles of the Quail (Table 5). Thus, the content of stearic fatty acid increased in the poultry red meat (groups III and IV) by 0.26% and 0.58% respectively.

However, the accumulation of myristic acid in these groups was less by 0.04% and 0.03% compared to the control. We found that linoleic acid content increased by 1.75%, 2.48% and 2.67% in groups I-III compared to the control group. We must note that the level of accumulation of γ -linolenic and α -linolenic fatty acids increased by 0.03% and 0.21% in group IV relative to the control. We also registered an increase in arachidonic fatty acid content by 0.39%, 2.39%, and by 1.78% in groups II-IV compared to the control. While the content of oleic and palmitoleic fatty acids was lower in all the groups, however, the highest rate of accumulation of dihomo- γ -linolenic, docosatetraenoic and docosahexaenoic fatty acids was observed in the femur muscles of Quails from group IV, they were higher by 0.18%, 0.38%, and 1.4%, respectively, compared to the control group. Therefore, the use of different doses of *Echinacea pallida* extract in Quail feeding had a positive effect on the amino acid, fatty acid, and chemical composition of the Quails meat.

Conclusion

The supplementation of the dry extract of *Echinacea pallida* increased the synthesis of essential amino acids in white Quail meat, namely lysine by 0.45%, histidine by 0.11%, arginine by 0.37%, valine by 0.24%, isoleucine by 0.16%, and phenylalanine by 0.08%. It was also increased the amount of essential amino acids by 1.81%. The herbal supplements application in Quail compound feed caused the increase of substitutable amino acids content in red meat, namely serine by 0.06%, proline by 0.12%, glycine by 0.03%. This was also caused the increase of the amount of substitutable amino acids by 2.28%.

The application of *Echinacea pallida* extract in Quails feeding caused the increase of fatty acids in the pectoral and femur muscles, namely linoleic acid by 5.41%, dihomo- γ -linolenic by 0.18%, arachidonic by 3.22% and linoleic by 2.48%, stearic by 0.26%, and arachidonic acid by 2.39%.

References

- Allen, P. (2003). Dietary supplementation with Echinacea and development of immunity to challenge infection with coccidian. *Parasitol. Res.*, V. 91, 74–78.
- Aituan, M. A., Wanyu, S. H. (2009). Effects of Echinacea purpurea extract on the immunological response to infectious bursal disease vaccine in broilers. Higher Education Press and Springer. Vol. 3(4), 452–456.
- Barnes, J., Anderson, L.A., Gibbons, S., Phillipson, J.D. (2005). Echinacea species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): a review of their chemistry, pharmacology and clinical properties. *Journal of Pharmacy and Pharmacology*, Vol. 57(8), 929–954.
- Bauer, R., Khan, J.A., Wagner, H. (1988). Analysis of *Echinacea pallida* and *E. angustifolia* Roots. *Planta Medica*, Vol. 54(5), 426–430.
- Bauer, R., Wagner, H. (1990). *Echinacea Handbuch für Ärzte, Apotheker und andere Naturwissen*. Stuttgart: Schaffler. 182. (in German)
- Bauer, R., Wagner, H. (1991). Echinacea species as nutritional potential immunostimulatory drugs. *Economic and Medicinal Plant Research*, Vol. 5, 253–321.
- Bohmer, M., Salisch, H. (2009). Echinacea purpurea as a potential immunostimulatory feed additive in laying hens and fattening pigs by intermittent application. *Livest. Sci*, 122, 81–85.
- Chudak, R.A., Poberezhets, Y.M., Vozniuk, O.I. (2018). Effect of *Echinacea pallida* dry extract on the liver composition of Quails. *Agrarian Science and Food Technology, VNAU*, 2(101), 81–89 (in Ukrainian)
- Chudak, R.A., Poberezhets, Y.M., Vozniuk, O.I., Dobronetska, V.O. (2019). Echinacea pallida extract effect on quails meat quality. *Ukrainian journal of ecology*, Vol 9(2), 151–155.
- Cross, D.E., Hillman, K. 2004. Antibacterial properties of phytochemicals in aromatic plants in poultry diets. *Poultry world*, 10, 19–23.
- Dehkordi, S., Fallah, V. (2011). Enhancement of broiler performance and immune response by Echinacea purpurea supplemented in diet. *Journal Home*, 10(54), 24–32.
- Diakonova, Ya. V. (2009). Pharmacogenetic study of *Echinacea pallida* Nutt: thesis abstract: spec. 15.00.02 Pharmaceutical chemistry and pharmacognosy. Kyiv (in Ukrainian)
- Grashorn, M. A. (2010). Use of phytobiotics in broiler nutrition – an alternative to in feed antibiotics. *Journal of Animal and Feed Sciences*, 19, 338–347.
- Ibatullin, I. I., Omelian, A. M., Sychov, M. Yu. (2017). Impact of different levels of arginine on zootechnical indices and slaughter characteristics of young Quails. *Ukrainian journal of ecology*, 7(1), 37–45. doi.org/10.15421/20174
- Ibatullin, I. I., Zhukorskyi, O. M., Bashchenko, I. (2017). Methodology and organization of scientific research in animal husbandry. Kyiv, Ahrar. Nauka (in Ukrainian).

- Magomedaliev, I.M., Nekrasov, R.V., Chabaev, M.G., Dzhavakhiya, V.V., Glagoleva, E.V., Matsyura, A. V. (2019). Use of different concentrations of enzymesporin probiotic in feeding of growing young pigs. *Ukrainian Journal of Ecology*, 9(4), 705-708. Doi: 10.15421/2019_152
- Mamchur, F. I., Zuzuk, B. M., Vasylyshyn, A. A. (1993). Chemical composition and pharmacological properties of plants of the genus *Echinacea* (Asteraceae). *Pharmaseutical Journal*, 2, 38–40. (in Ukrainian)
- Nazir, Z. and Grashorn, M. (2010). Effect of *Echinacea purpurea* and *Nigella sativa* supplementation on broiler performance, carcass and meat quality. *Journal of Animal and Feed Sciences*, 9, 94–104.
- Podolian, Ju. N. (2017). Effect of probiotics on the chemical, mineral, and amino acid composition of broiler chicken meat. *Ukrainian journal of ecology*, 7(1), 61–65. DOI: <http://dx.doi.org/10.15421/20178>
- Podolian, Yu. M. (2016). Influence of probiotic on productivity of broiler chickens. *Biological journal of the Melitopol State Pedagogical University named after Bogdan Khmelnytsky*, 6(3), 141-148. DOI: <http://dx.doi.org/10.15421/201680>. (in Ukrainian)
- Pospelov, S. V., Samorodov, V. N., Dyakonova, Ya. V. (2007). Ontomorphology and phytochemistry of *Echinacea pallida* (Nutt.) introduced in Ukraine. *Biological diversity. Plant Introduction, Proceedings of the Fourth Int. conf. SPb.*, 500–501. (in Russian)
- Samorodov, V. N., Pospelov, S. V. (2013). The results of the study and selection of representatives of the genus *Echinacea* Moench in the Poltava State Agrarian Academy. *Proceedings of the international scientific conference Innovative approaches to the Echinacea research. Poltava*, 89–99. (in Russian).
-

Citation:

Chudak, R.A., Ushakov, V.M., Poberezhets, Y.M., Lotka, H.I., Polishchuk, T.V., Kazmiruk, L.V. (2020). Effect of *Echinacea pallida* supplementation on the amino acid and fatty acid composition of Pharaoh Quail meat. *Ukrainian Journal of Ecology*, 10(2), 302-307.



This work is licensed under a Creative Commons Attribution 4.0. License
