
НАУКОВІ ГОРИЗОНТИ



Засновник, редакція, видавець
ПОЛІСЬКИЙ НАЦІОНАЛЬНИЙ УНІВЕРСИТЕТ

**Свідоцтво про державну реєстрацію
Серія KB № 24756-14696 ПР від 26.02.2021 р.**

Науковий журнал включено до категорії Б Переліку наукових фахових видань України, в яких можуть публікуватися результати дисертаційних робіт на здобуття наукових ступенів доктора і кандидата ветеринарних, економічних, сільськогосподарських та технічних наук зі спеціальностей – 051, 071, 072, 073, 075, 076, 101, 133, 183, 201, 202, 203, 204, 205, 206, 208, 211, 281, 292 (наказ МОН України № 1643 від 28.12.2019 р., наказ МОН України № 409 від 17.03.2020 р.).

Журнал включено до міжнародних наукометричних баз і каталогів наукових видань: Index Copernicus; Directory of Open Access Journals (DOAJ); Open Academic Journals Index (OAJI); Google Scholar; Crossref; Національна бібліотека України імені В.І. Вернадського.

Друкується за рішенням Вченої ради Поліського національного університету, протокол № 13 від 23.06.2021 р.

ISSN: 2663-2144
e-ISSN: 2709-8877

Підписано до друку 23.06.2021 р.
Формат 210×297. Ум. друк. арк. 14,6
Наклад 90 примірників
© Поліський національний університет, 2021

SCIENTIFIC HORIZONS



Founder, Editorial and Publisher
POLISSIA NATIONAL UNIVERSITY

Certificate of state registration
KV No. 24756-14696 PR of February 26, 2021.

The scientific journal is included in category B of the List of scientific professional periodicals of Ukraine. It enables publishing the thesis results for Doctor and Candidate degrees in economic agricultural, technical and veterinary sciences (Order of the Ministry of Education and Science of Ukraine No 1643 of December 28, 2019; Order of the Ministry of Education and Science of Ukraine No 409 of March 18, 2020). It comprises the following specialties – 051, 071, 072, 073, 075, 076, 101, 133, 183, 201, 202, 203, 204, 205, 206, 208, 211, 281, 292.

The journal is included in the international scientific databases and catalogs of scientific publications: Index Copernicus; Directory of Open Access Journals (DOAJ); Open Academic Journals Index (OAJI); Google Scholar; Crossref; National Library of Ukraine named after V. I. Vernadskiy.

Recommended for publication by the decision of the Academic Council Polissia National University Minutes No. 13 of 23.06.2021.

ISSN: 2663-2144
e-ISSN: 2709-8877

Signed for publication 23.06.2021
Format 210×297. Mind. print. ark. 14.6.
Circulation 90 copies
© Polissia National University, 2021

НАУКОВІ ГОРИЗОНТИ

Том 24, № 2 2021

НАУКОВИЙ ЖУРНАЛ
Засновано 12 березня 1998 р.
Періодичність випуску: чотири рази на рік

Редакційна колегія

Головний редактор

Олег Васильович Скидан

ректор, д-р екон. наук, професор, Поліський національний університет, Україна

Заступник головного редактора

Л. Романчук

д-р с.-г. наук, Поліський національний університет, Україна

Члени редакційної колегії

Ю. Раманаускас

д-р наук, Клайпедський університет, Литва

С. Кухарець

д-р техн. наук, Поліський національний університет, Україна

І. Грабар

д-р техн. наук, Поліський національний університет, Україна

Я. Ярош

д-р техн. наук, Поліський національний університет, Україна

Т. Федонюк

д-р с.-г. наук, Поліський національний університет, Україна

Н. Сорока

д-р вет. наук, Національний університет біоресурсів і природокористування України, Україна

Р. Ставецька

д-р с.-г. наук, Білоцерківський національний аграрний університет, Україна

В. Шлапак

д-р с.-г. наук, Уманський національний університет садівництва, Україна

І. Левкович

д-р наук, Лейбніцький інститут розвитку сільського господарства у країнах з перехідною економікою, Німеччина

А. Зимароєва

канд. біол. наук, Поліський національний університет, Україна

П. Чумак

канд. с.-г. наук, Поліський національний університет, Україна

Л. Чижевська

д-р екон. наук, Державний університет «Житомирська політехніка», Україна

В. Данкевич

д-р екон. наук, Поліський національний університет, Україна

Т. Асанбаєв

канд. с.-г. наук, Павлодарський державний університет ім. С. Торайгирова, Республіка Казахстан

Ю. Джуянді

д-р філософ. наук, Паджанджаранський університет, Індонезія

О. Патапас

д-р філософ. наук, Університет Миколаса Ромеріса, Литва

Я.-У. Сандал

ректор, професор, д-р філол. наук, Інститут Доктора Яна-У. Сандала, Норвегія

Е. Сараускіс

професор, Інститут сільськогосподарської техніки та безпеки університету Вітовта Великого, Литва

Р. Пукала

канд. екон. наук, Державний Техніко-Економічний Університет ім. кс. Броніслава Маркевича в Ярославі, Польща

Е. Алієв

д-р техн. наук., Інститут олійних культур Національної академії аграрних наук України, Україна

О. Галатюк

д-р вет. наук, Поліський національний університет, Україна

В. Гамаюнова

д-р с.-г. наук, Миколаївський національний аграрний університет, Україна

Л. Горальський

д-р вет. наук, Поліський національний університет, Україна

С. Гуральська

д-р вет. наук, Поліський національний університет, Україна

Є. Гулієва

канд. хім. наук, Національна академія наук Азербайджану, Азербайджан

Д. Дерев'янок

д-р техн. наук, Поліський національний університет, Україна

Ж.-П. Вандерлінден

д-р екон. наук, директор дослідного центру CEARC, Франція

О. Жуков

д-р біол. наук, Мелітопольський державний педагогічний університет імені Богдана Хмельницького, Україна

І. Іванова	канд. с.-г. наук, Таврійський державний агротехнологічний університет ім. Д. Моторного, Україна
М. Ключевич	д-р с.-г. наук, Поліський національний університет, Україна
О. Кочук-Ященко	канд. с.-г. наук, Поліський національний університет, Україна
І. Кравчук	д-р екон. наук, Поліський національний університет, Україна
Д. Кучер	канд. с.-г. наук, Поліський національний університет, Україна
І. Лігоміна	д-р вет. наук, Поліський національний університет, Україна
Д. Лісогурська	канд. с.-г. наук, Поліський національний університет, Україна
Ф. Марков	канд. с.-г. наук, Поліський національний університет, Україна
О. Марковська	д-р с.-г. наук, Херсонський державний аграрно-економічний університет, Україна
А. Михайлов	д-р екон. наук, Сумський національний аграрний університет, Україна
В. Мойсієнко	д-р с.-г. наук, Поліський національний університет, Україна
В. Мороз	канд. біол. наук, Поліський національний університет, Україна
Ю. Мороз	д-р екон. наук, Поліський національний університет, Україна
В. Москалець	д-р с.-г. наук, Інститут садівництва НААН, Україна
Ю. Никитюк	д-р екон. наук, Поліський національний університет, Україна
В. Пазич	канд. с.-г. наук, Поліський національний університет, Україна
В. Соколюк	д-р вет. наук, Поліський національний університет, Україна
С. Танірбергенов	вчений секретар, Казахський науково-дослідний інститут ґрунтознавства і агрохімії ім. У.У. Успанова, Республіка Казахстан
Л. Тарасович	канд. екон. наук, Поліський національний університет, Україна
Т. Тимошук	канд. с.-г. наук, Поліський національний університет, Україна
Н. Цивенкова	канд. техн. наук, Поліський національний університет, Україна
М. Швець	канд. біол. наук, Поліський національний університет, Україна
С. Щетина	канд. с.-г. наук, Уманський національний університет садівництва, Україна

SCIENTIFIC HORIZONS

Vol. 24, No. 2 2021

SCIENTIFIC JOURNAL
Year of establishment: since March 1998.
Publication frequency: four times a year

Editorial Board

Editor-in-Chief

O. V. Skydan

Rector, Full Doctor in Economic Sciences, Professor, Polissia National University, Ukraine

Deputies Editor-in-Chief

L. Romanchuk

Full Doctor in Agricultural Sciences, Polissia National University, Ukraine

Editorial board members

J. Ramanauskas

Dr. Habil., Klaipeda University, Lithuania

S. Kukharets

Full Doctor in Engineering Sciences, Polissia National University, Ukraine

I. Grabar

Full Doctor in Engineering Sciences, Polissia National University, Ukraine

Y. Yarosh

Full Doctor in Engineering Sciences, Polissia National University, Ukraine

T. Fedoniuk

Full Doctor in Agricultural Sciences, Polissia National University, Ukraine

N. Soroka

Full Doctor in Veterinary Sciences, National University of Life and Environmental Sciences of Ukraine, Ukraine

R. Stavetska

Full Doctor in Agricultural Sciences, Bila Tserkva National Agrarian University, Ukraine

V. Shlapak

Full Doctor in Agricultural Sciences, Uman National University of Horticulture, Ukraine

I. Levkovych

Dr. Habil., Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Germany

A. Zymarioieva

PhD in Biological Sciences, Polissia National University, Ukraine

P. Chumak

PhD in Agricultural Sciences, Polissia National University, Ukraine

L. Chyzhevska

Full Doctor in Economic Sciences, Zhytomyr Polytechnic State University, Ukraine

V. Dankevych

Full Doctor in Economic Sciences, Polissia National University, Ukraine

T. Assanbayev

PhD in Agricultural Sciences, Pavlodar State University named after S. Toraiyrov, Republic of Kazakhstan

Yu. Djuyandi

Full Doctor in Philosophical Sciences, Padjadjaran University, Indonesia

A. Patapas

Full Doctor in Philosophical Sciences, Mykolas Romeris University, Lithuania

Ja.-U. Sandal

Rector, Professor, Full Doctor in Philological Sciences, Jan-U. Sandal Institute, Norway

E. Sarauskis

Professor, Institute of Agricultural Engineering and Safety of Vytautas Magnus university (VMU), Lithuania

R. Pukala

PhD in Economic Sciences, The Bronisław Markiewicz State University of Technology and Economics in Jarosław, Poland

E. Aliiev

Full Doctor in Engineering Sciences, Institute of Oilseed Crops of the National Academy of Agricultural Sciences of Ukraine, Ukraine

O. Galatiuk

Full Doctor in Veterinary Sciences, Polissia National University, Ukraine

V. Hamaiunova

Full Doctor in Agricultural Sciences, Mykolayiv National Agrarian University, Ukraine

L. Goralskiy

Full Doctor in Veterinary Sciences, Polissia National University, Ukraine

S. Guralaska

Full Doctor in Veterinary Sciences, Polissia National University, Ukraine

E. Guliyeva

PhD in Chemical Sciences, National Academy of Sciences of Azerbaijan, Azerbaijan

D. Derevjanko	Full Doctor in Engineering Sciences, Polissia National University, Ukraine
J.-P. Vanderlinden	Full Doctor in Economic Sciences, Director of the CEARC Research Center, France
O. Zhukov	Full Doctor in Biological Sciences, Bogdan Khmelnytsky Melitopol State Pedagogical University, Ukraine
I. Ivanova	PhD in Agricultural Sciences, Dmytro Motornyi Tavria State Agrotechnological University, Ukraine
M. Kliuchevich	Full Doctor in Agricultural Sciences, Polissia National University, Ukraine
O. Kochuk-Yashchenko	PhD in Agricultural Sciences, Polissia National University, Ukraine
I. Kravchuk	Full Doctor in Economic Sciences, Polissia National University, Ukraine
D. Kucher	PhD in Agricultural Sciences, Polissia National University, Ukraine
I. Ligomina	Full Doctor in Veterinary Sciences, Polissia National University, Ukraine
D. Lisohurska	PhD in Agricultural Sciences, Polissia National University, Ukraine
F. Markov	PhD in Agricultural Sciences, Polissia National University, Ukraine
O. Markovska	Full Doctor in Agricultural Sciences, Kherson State Agrarian and Economic University, Ukraine
A. Mykhailov	Full Doctor in Economic Sciences, Sumy National Agrarian University, Ukraine
V. Moysiienko	Full Doctor in Agricultural Sciences, Polissia National University, Ukraine
V. Moroz	PhD in Biological Sciences, Polissia National University, Ukraine
J. Moroz	Full Doctor in Economic Sciences, Polissia National University, Ukraine
V. Moskalets	Full Doctor in Agricultural Sciences, Institute of Horticulture of the National Academy of Agrarian Sciences, Ukraine
Yu. Nykytiuk	Full Doctor in Economic Sciences, Polissia National University, Ukraine
V. Pazyh	PhD in Agricultural Sciences, Polissia National University, Ukraine
V. Sokolyuk	Full Doctor in Veterinary Sciences, Polissia National University, Ukraine
S. Tanirbergenov	Scientific Secretary, U.U. Uspanov Kazakh Research Institute of Soil Science and Agrochemistry, Republic of Kazakhstan
L. Tarasovych	PhD in Economic Sciences, Polissia National University, Ukraine
T. Tymoshchuk	PhD in Agricultural Sciences, Polissia National University, Ukraine
N. Tsyvenkova	PhD in Engineering Sciences, Polissia National University, Ukraine
M. Shvets	PhD in Biological Sciences, Polissia National University, Ukraine
S. Shchetyna	PhD in Agricultural Sciences, Uman National University of Horticulture, Ukraine

ЗМІСТ

Б. А. Шелудченко, І. П. Слюсаренко, О. Б. Плужніков, В. О. Шубенко, В. Р. Білецький, В. М. Боровський АНАЛІТИЧНИЙ КРИТЕРІЙ МІЦНОСТІ ЗВ'ЯЗАНО-ДИСПЕРСНИХ ГЕЛЕЙ ПРИ ТРАНСПОРТУВАННІ ЇХ У ТРУБОПРОВОДАХ.....	9
Е. Б. Алієв, Р. Д. Малегін, В. В. Івлєв, О. Ю. Алієва СИМУЛЯЦІЯ ПРОЦЕСУ КАВІТАЦІЙНОЇ ОБРОБКИ РІДКИХ КОРМІВ.....	16
С. В. Ребко, О. В. Козел, І. В. Кімейчук, В. Ю. Юхновський ПОРІВНЯЛЬНА ОЦІНКА ДЕЯКИХ ФІЗИКО-МЕХАНІЧНИХ ВЛАСТИВОСТЕЙ ДЕРЕВИНИ РІЗНИХ КЛІМАТИПІВ СОСНИ ЗВИЧАЙНОЇ.....	27
О. Г. Василевський, І. С. Нейко, Ю. А. Єлісавенко, М. В. Матусяк ХАРАКТЕРИСТИКА ДУБОВИХ ЛІСІВ ПРИРОДНОГО ПОХОДЖЕННЯ ДП «ХМІЛЬНИЦЬКЕ ЛГ» ТА ОСОБЛИВОСТІ ЗАПРОВАДЖЕННЯ ЗАХОДІВ ЩОДО ЇХ ВІДТВОРЕННЯ.....	37
А. Л. Шуляр, А. Л. Шуляр, В. Ф. Андрійчук, С. П. Омелькович, В. П. Ткачук, В. В. Латка ПОРІВНЯЛЬНА ОЦІНКА ПРОДУКТИВНИХ ОЗНАК СВИНЕЙ РІЗНИХ ПОРІД І ПОРОДНИХ ПОЄДНАНЬ.....	47
М. С. Коваленко, Д. Д. Білий ПРОГНОСТИЧНЕ ЗНАЧЕННЯ СУДИННОЇ ІНВАЗІЇ ЗА ПУХЛИН МОЛОЧНОЇ ЗАЛОЗИ У СУК (ПІЛОТНЕ ДОСЛІДЖЕННЯ).....	54
С. В. Маслійов, Н. О. Коржова ФОТОСИНТЕТИЧНИЙ ПОТЕНЦІАЛ РОСЛИН ЯЧМЕНЮ ЯРОГО В УМОВАХ ЗОНИ СТЕПУ.....	62
О. П. Ткачук БІОЛОГІЧНІ ОСОБЛИВОСТІ ПОШИРЕННЯ КОРЕНЕВИХ СИСТЕМ БОБОВИХ БАГАТОРІЧНИХ ТРАВ В УМОВАХ ЗМІНИ КЛІМАТУ.....	69
В. Г. Дідора, М. М. Ключевич ПРОДУКТИВНІСТЬ СОЇ ЗАЛЕЖНО ВІД ЕЛЕМЕНТІВ ОРГАНІЧНОЇ ТЕХНОЛОГІЇ ВИРОЩУВАННЯ В КОРОТКОРОТАЦІЙНІЙ СІВОЗМІНІ ПОЛІССЯ УКРАЇНИ.....	77
Ю. Ю. Чуприна КЛАСТЕРНИЙ АНАЛІЗ ЗРАЗКІВ <i>TRITICUM L.</i> РІЗНОГО ЕКОЛОГО-ГЕОГРАФІЧНОГО ПОХОДЖЕННЯ.....	84
Р. В. Яковенко, П. Г. Копитко, В. М. Пелехатий ВМІСТ ХЛОРОФІЛУ ТА ЕЛЕМЕНТІВ ЖИВЛЕННЯ В ЛИСТІ ЯБЛУНІ ЗАЛЕЖНО ВІД ДОВГОТРИВАЛОГО УДОБРЕННЯ.....	93
А. А. Коробко ОЦІНКА СОРТІВ КВІТНИКОВО-ДЕКОРАТИВНИХ РОСЛИН ВИДУ <i>NEMESIA VENT</i> В УМОВАХ ПОДІЛЛЯ.....	99
Т. І. Дзюба ІНФОРМАЦІЙНА МОДЕЛЬ УДОСКОНАЛЕННЯ ОБЛІКОВО-АНАЛІТИЧНОГО ЗАБЕЗПЕЧЕННЯ УПРАВЛІННЯ ЕКОНОМІЧНИМ ПОТЕНЦІАЛОМ.....	108

CONTENTS

B. Sheludchenko, I. Slusarenko, O. Pluzhnikov, V. Shubenko, V. Biletsky, V. Borovskyi ANALYTICAL CRITERION OF STRENGTH CONNECTED-DISPERSED GELS WHEN TRANSPORTING THEM IN PIPELINES.....	9
E. Aliiev, R. Maliehin, V. Ivliev, O. Aliieva SIMULATION OF THE PROCESS OF CAVITATION TREATMENT OF LIQUID FEED.....	16
S. Rabko, A. Kozel, I. Kimeichuk, V. Yukhnovskyi COMPARATIVE ASSESSMENT OF SOME PHYSICAL AND MECHANICAL PROPERTIES OF WOOD OF DIFFERENT SCOTS PINE CLIMATYPES.....	27
O. Vasylevskyi, I. Neyko, Yu. Yelisavenko, M. Matusia CHARACTERISTICS OF NATURAL OAK FORESTS OF IN SE "KHMILNYTSKE LISOVE HOSPODARSTVO" AND IMPLEMENTATION OF MEASURES FOR THEIR REGENERATION	37
A. Shuliar, A. Shuliar, V. Andriichuk, S. Omelkovych, V. Tkachuk, V. Latka COMPARATIVE ASSESSMENT OF ECONOMIC TRAITS OF PIGS OF DIFFERENT BREEDS AND BREED COMBINATIONS.....	47
M. Kovalenko, D. Bilyi PROGNOSTIC VALUE OF VASCULAR INVASION IN BREAST TUMOURS IN SHE-DOGS (PILOT STUDY)	54
S. Masliiov, N. Korzhova PHOTOSYNTHETIC POTENTIAL OF SPRING BARLEY PLANTS IN THE STEPPE ZONE	62
O. Tkachuk BIOLOGICAL FEATURES OF THE DISTRIBUTION OF ROOT SYSTEMS OF PERENNIAL LEGUME GRASSES IN THE CONTEXT OF CLIMATE CHANGE	69
V. Didora, M. Kluchevych SOYBEAN PRODUCTIVITY DEPENDING ON THE ELEMENTS OF ORGANIC CULTIVATION TECHNOLOGY IN THE SHORT-TERM CROP ROTATION OF UKRAINIAN POLISSIA.....	77
Yu. Chuprina CLUSTER ANALYSIS OF <i>TRITICUM L.</i> SAMPLES OF DIFFERENT ECOLOGICAL AND GEOGRAPHICAL ORIGIN	84
R. Yakovenko, P. Kopytko, V. Pelekhatyi THE CONTENT OF CHLOROPHYLL AND NUTRIENTS IN APPLE LEAVES DEPENDING ON LONG-TERM FERTILISER.....	93
A. Korobko EVALUATION OF FLOWER DECORATIVE PLANTS CULTIVARS OF <i>NEMESIA VENT</i> IN THE CONDITIONS OF PODILLIA	99
T. Dziuba INFORMATION MODEL FOR IMPROVING ACCOUNTING AND ANALYTICAL SUPPORT FOR ECONOMIC POTENTIAL MANAGEMENT	108

SCIENTIFIC HORIZONS

Journal homepage: <https://sciencehorizon.com.ua>

Scientific Horizons, 24(2), 69-76



UDC 574/577/.58.036.1

DOI: 10.48077/scihor.24(2).2021.69-76

BIOLOGICAL FEATURES OF THE DISTRIBUTION OF ROOT SYSTEMS OF PERENNIAL LEGUME GRASSES IN THE CONTEXT OF CLIMATE CHANGE

Oleksandr Tkachuk*

Vinnytsia National Agrarian University
21008, 3 Soniachna Str., Vinnytsia, Ukraine

Article's History:

Received: 01.03.2021

Revised: 10.04.2021

Accepted: 16.05.2021

Suggested Citation:

Tkachuk, O. (2021). Biological features of the distribution of root systems of perennial legume grasses in the context of climate change. *Scientific Horizons*, 24(2), 69-76.

Abstract. The key factor in the choice of perennial legume grasses in the face of climate change is their drought resistance, because the development of roots directly affects this property, as well as determines their fertility. The purpose of the study was to establish what morphological and biological characteristics of grass roots ensure economic characteristics in the context of climate change. Methods of observation, comparison, and field experiment were used. The study found that the roots of alfalfa have the greatest depth in the soil, spread in the horizontal direction, the thickness of the root neck and lateral roots, which ensures plasticity, durability, and productivity of its crops. The roots of white melilot have the greatest depth in the soil, the thickness of the root neck, central root and lateral roots, their spread in the horizontal direction, which affects the drought and frost resistance of crops. The roots of birdsfoot trefoil are distinguished by the greatest number of renewal buds on the root neck, the smallest depth, and the thickness of the root neck, which affects productive longevity and the possibility of growth in poor and acidic soils. The roots of eastern galega have the greatest depth of the main plant, distribution in the horizontal direction, the thickness of the central root, which affects productive longevity and high biological plasticity. The roots of Hungarian sainfoin are distinguished by the greatest distribution in the horizontal direction and the deepest placement of the main branching, which affects high biological resistance to adverse growing conditions. It was also found out that the roots of meadow clover have the smallest spread radius in the horizontal plane, the thickness of the central and lateral roots, which determines the possibility of its cultivation in the field crop rotation. These findings would facilitate the selection of perennial legume grasses to achieve their full potential in the face of climate change

Keywords: plants, roots, underground development, morphology, years of vegetation



Copyright © The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (<https://creativecommons.org/licenses/by/4.0/>)

*Corresponding author

INTRODUCTION

One of the key factors when choosing perennial legume grasses in the face of climate change is their adaptability to periodic lack of moisture during the growing season, which depends on the development of the root system of grasses. At the same time, placing crops of perennial legume grasses on poor and acidic soils limits the growth of their roots and reduces drought resistance [1]. It is known that the processes that occur in the vegetative organs of plants have a direct connection with the function of their underground part. The larger the volume and length of the root system, the greater the aboveground mass of the plant. A well-developed root system contributes to the efficient use of nutrients and moisture from the soil [2].

In arid conditions, in the year of sowing with clean cultivation, the roots of alfalfa penetrate to a depth of 120-140 cm. With sufficient moisture supply, the root system does not need to penetrate deeply. In general, the root of alfalfa penetrates to a depth of 10 m or more [3]. The root system of meadow clover penetrates the soil from 20 cm per year of sowing and up to 150-200 cm in general. Hungarian sainfoin has a well-developed root system that penetrates up to 150-200 cm [4]. Birdsfoot trefoil is less resistant to drought than alfalfa. Its root system is well branched and penetrates up to 1.5 m deep and is well adapted to acidic and poor soils [5]. Eastern galega is also drought-resistant but develops well in soils with high water retention capacity [1]. White melilot has a well-developed root system, which determines its drought resistance. In white melilot, the root is thickened in the upper part, with well-developed lateral roots, penetrates deeply into the soil, reaching the subsoil [6].

The issue of the distribution of root systems of traditional perennial legume grasses in the context of climate change has been studied in the works of many scientists. In particular, G.P. Kvitko investigated the influence of the development of root systems on soil fertility indicators [5]; P.S. Makarenko studied the distribution of plant roots in grass mixtures [6]; O.L. Kirilesko considered the development of root systems of perennial legume grasses on sloping lands [7]; I. Zabarna examined the morphology of meadow clover roots [8]. K. Schnidtko, R. Rauber analysed the influence of root development on the intensity of their absorption of heavy metals from the soil [9]; P.K. Ghosh and T.K. Maiti studied the relationship between the root systems of perennial legumes and their nodule bacteria [10]; M. Marczak, A. Mazur, P. Koper, K. Zebracki and A. Skorupska investigated the symbiotic productivity of grass roots [11]; G.G. Stevens, M.A. Perez-Fernandez, R.J.L. Morcillo, A. Kleinert, P. Hills, D.J. Brand, E.T. Steenkamp and A.J. Valentine studied the peculiarities of growth and development of the roots of perennial legume grasses in conditions of nutrient deficiency in the soil [12]; T. Suzuki, N. Takeda, H. Nishida, M. Hoshino, M. Ito,

F. Misawa, Y. Handa, K. Miura and M. Kawaguchi studied the development of root systems of perennial legume grasses in the absence of symbiotic bacteria in the soil [13]; C. Jacob, B. Carrasco, A.R. Schwember analysed the achievements of selection and biotechnology of perennial legumes [14]; M.A. Adams, N. Buchmann, J.I. Sprent, T.N. Buckley and T.L. Turnbull examined the effects of moisture and nitrogen on the spread of root systems of grass [15]. For the most part, they indicate a powerful development of the root system. At the same time, the influence of climate change conditions on the morphological features of the root systems of perennial legumes and their adaptation to adverse environmental conditions has been rather neglected.

Alfalfa is a traditional forest-steppe crop, has been grown in this region for a long time, is well adapted to the soil and climatic factors and changes in growing conditions, and characterised by plasticity and durability. However, the use of other types of perennial legume grasses should be complementary to alfalfa, using their biological and economic advantages in local soil and climatic conditions. Thus, Hungarian sainfoin is characterised by high biological resistance to adverse growing conditions; white melilot has good drought and winter hardiness. Birdsfoot trefoil is characterised by productive longevity and the ability to grow on poor and acidic soils. Eastern galega is characterised by productive longevity and high biological plasticity [16].

Therefore, the *purpose* of this study was to establish what morphological and biological characteristics of the root systems of perennial legume grasses ensure their ecological and economic features in the context of climate change.

MATERIALS AND METHODS

Field research was conducted during 2013-2017 at the research farm "Agronomichne" of the Vinnytsia National Agrarian University. The experimental site was located on gray podzolic medium loamy soil. Agrochemical composition of the soil: humus content – 2.0%, hydrolysed nitrogen (according to Kornfeld) – 133 mg/kg of soil, mobile phosphorus (according to Chirikov) – 390 mg/kg of soil, mobile potassium (according to Chirikov) – 64 mg/kg of soil, hydrolytic acidity – 2.53 mg-EQ/100 g of soil, reaction of pH salt of soil solution – 6.0.

Studies were conducted with the following types of perennial legume grasses: alfalfa (*Medicago sativa* L.), red clover (*Trifolium pratense* L.), Hungarian sainfoin (*Onobrychis arenaria* Kit.), white melilot (*Melilotus albus* L.), birdsfoot trefoil (*Lotus corniculatus* L.) and eastern galega (*Galega orientalis* Lam.). Grasses were sown in early spring using a clean cultivation way using a post-emergence herbicide. The green mass was harvested at the beginning of the flowering phase. Determination of the distribution features of the roots of perennial legume grasses in the soil was carried out by dry

excavation [17; 18]. Quantitative parameters of root systems were determined using a measuring tape and caliper in four repetitions.

During the 2013 calendar year, 652 mm of precipitation fell, which is 18 mm more than the long-term average (634 mm), with an average annual temperature of 9.0°C, which is 2.0°C more than normal (7.0°C). The growing season began in the first ten days of April and lasted until the end of the second ten days of November. During the growing season, the amount of precipitation was 429 mm. In 2014, precipitation was 550 mm, which was 87% of the long-term average. The average annual temperature was 8.6°C, which is 1.6°C higher than normal. The growing season began in the first ten days of March and lasted until the end of the first ten days of November. During the growing season, the amount of precipitation was 442 mm.

In 2015, 368 mm of precipitation fell, which was 58% of the long-term average. The average annual temperature was 9.3°C, which is 2.3°C higher than normal. The growing season began in the end of March and lasted until the second ten days of November. During the growing season, 235 mm of precipitation fell. In 2016, the average annual temperature was

9.0°C, which is 2°C higher than normal. The amount of precipitation during the year was 469 mm, which was 26% less than normal. The growing season began in early April and lasted until the end of September. 2017 was characterised by an average annual temperature of 9.1°C, which was 2.1°C higher than normal. The amount of precipitation for the year was 503 mm, which corresponds to 80% of the long-term norm.

Thus, the most favourable growing conditions, taking into account the temperature regime and humidity level, were in 2014, which corresponds to the second year of vegetation of perennial legume grasses. The least favourable growing conditions were typical observed in 2016, when the grasses were growing for the fourth year.

RESULTS AND DISCUSSION

Features of initial aboveground and underground growth of perennial legume grasses

Observations of the germination of root systems of perennial legume grasses have shown that the roots of alfalfa and white melilot begin to germinate most quickly – on the 2nd-3rd day after sowing, later – red clover, Hungarian sainfoin, and at the latest – birdsfoot trefoil (Table 1).

Table 1. Dynamics of early growth of underground and aboveground parts of perennial legume grasses, M±m

Perennial legume grasses	Height, cm depending on the day after sowing				
	5		10		15
	underground growth	aboveground growth	underground growth	aboveground growth	underground growth
Alfalfa	2.1±0.2	0.6±0.2	2.5±0.2	1.1±0.2	4.2±1.0
Red clover	2.2±0.1	0.2±0.1	2.6±0.2	0.6±0.1	2.9±0.4
Hungarian sainfoin	2.7±0.3	0.5±0.2	3.7±0.3	3.5±0.3	7.1±2.1
White melilot	1.7±0.2	0.3±0.1	2.2±0.2	0.8±0.1	3.3±0.9
Birdsfoot trefoil	0.3±0.1	0.1±0.05	0.8±0.1	0.4±0.1	2.5±1.0
Eastern galega	0.6±0.1	0.2±0.05	1.3±0.2	1.5±0.2	4.7±1.6

Source: compiled based on the author's own research

The results of studies showed that on the fifth day after sowing, the length of the roots of perennial legume grasses was 0.3-2.7 cm. Hungarian sainfoin had the greatest root length, which was 89% longer than birdsfoot trefoil, which crops had the smallest root length.

On the 10th day after sowing, the root length of perennial legume grasses was 0.8-3.7 cm. It remained the largest in plants of Hungarian sainfoin, which was 78.4% more than in plants of birdsfoot trefoil, which had the smallest root length. Starting from the 10th day, the growth of aboveground parts of legumes and perennial grasses begins. The ratio between the length of roots and aboveground seedlings in most plants was 6.5-8.0, in alfalfa – 4.2, and in red clover – 13.0 with a predominance of underground parts.

15 days after sowing, the length of the roots was 2.5-7.1 cm, and aboveground seedlings – 0.4-3.5 cm. In all indicators, Hungarian sainfoin plants prevailed. The ratio between the length of the underground and aboveground parts of plants decreased to 2.0-6.3. At this time, the root system grew more intensively in comparison with the aboveground part in red clover, birdsfoot trefoil and white melilot, and the aboveground part – in Hungarian sainfoin, eastern galega and alfalfa. The growth rate of the roots of perennial legume grasses for the first 5 days was highest in Hungarian sainfoin – 0.54 cm/day, while in birdsfoot trefoil – 0.06 cm/day.

During the second five-day period, the rate of root growth was 0.08-0.20 cm/day. It was the highest in the Hungarian sainfoin plants, and the smallest – in alfalfa

and red clover. During this period, the intensity of root growth decreases in all types of legume grasses, with the exception of birdsfoot trefoil and eastern galega. The greatest decrease in growth was observed in alfalfa and Meadow Clover – by 5.5-5.3 times, compared to the first five-day period. This is conditioned by the beginning of aboveground growth. Plants of birdsfoot trefoil and eastern galega increased the average daily growth, compared to the first period, by 1.7 and 1.2 times – up to 0.14-0.10 cm/day and had, respectively, the smallest growth of the aboveground part of the plant, compared to other plants.

In the third observation period, the average daily growth of root systems of plants was 0.22-0.68 cm/day, except for red clover, where this indicator was 0.06 cm/day. The largest gains were observed in Hungarian sainfoin and eastern galega. The intensity of root growth at this time increased by 2.2-4.9 times compared to the second five-day period, especially in the eastern galega.

The highest average daily root growth in the third observation period was noted in eastern galega, birdsfoot trefoil and Hungarian sainfoin. The intensity of their growth in the eastern galega and birdsfoot trefoil increased from the first to the third periods, while in the Hungarian sainfoin it decreased in the second period compared to the first. Alfalfa, red clover and white melilot had the greatest root gains in the first observation period.

Morphological and biological features of root system development

The intensity of growth and development of root systems of perennial legume grasses depends on the value of symbiotic nitrogen fixation of nodule bacteria, the intensity of absorption of nutrients and water from

the soil, the size of the formed aboveground crop. From an agroecological standpoint, the size of the root system determines the amount of organic matter that will return to the soil and replenish the supply of nutrients, and it also affects the agroecological resistance of the agrocenosis of legumes to unfavorable abiotic and biotic factors. The complex of morphological features of the distribution of plant root systems includes such parameters as the spread of roots in depth, the horizontal radius of root growth from the stem, and the depth of the main mass of branching of the plant root system.

The greatest depth of penetration of the root system into the soil thickness in the first year of vegetation was reached by alfalfa and white melilot, respectively, 83 and 82 cm, while the root system of birdsfoot trefoil reached a depth of 44 cm. The radius of horizontal root distribution, relative to the central root, was the largest in alfalfa plants – 25 cm, and the smallest – in red clover – 9 cm. The biggest depth of the main mass of branching of the root system was in eastern galega plants – 28 cm and Hungarian sainfoin – 23 cm, and the smallest – in alfalfa – 15 cm. A special feature of the root system of white melilot is its extremely high fragility, which may determine the biennial nature of the plant (Fig. 1).

Eastern galega, in addition to the root system itself, which is characterised by strong fibrosity, which determines its durability, also has root shoots, which are autonomous centres of emergence of individual plants. At the end of the first year of vegetation, the eastern galega plant forms up to 7-9 shoots. The maximum radius of their penetration from the central root is 25 cm, and the thickness is 3 mm. On such a shoot there can be up to 7 buds.




Figure 1. Root system of eastern galega (A), white melilot (B), alfalfa (C), Hungarian sainfoin (D) on the first year of growth

The main organ for restoring the vegetation of legume grasses and accumulating spare nutrients is the root neck. The productivity and durability of the herbage depends on the features of its development.

The greatest thickness of the root neck in the year of sowing was observed in white melilot – 12 mm and alfalfa – 11 mm, and the smallest – in eastern galega – 5 mm and red clover – 6 mm (Table 2).

Table 2. Morphology of root systems of perennial legumes depending on the growing season, $M \pm m$

Perennial legume grasses	Years of growth	Depth of root penetration, cm	Root system radius, cm	Depth of the main mass of branching, cm	Thickness, mm			Number of buds on the root neck, pcs.
					root neck	central root	side roots	
Alfalfa	1	83±3	25±3	15±1	11±1	4±1	1.5±0.2	12±3
	2	258±9*	28±3	26±2	18±2	5±1	2.0±0.2	25±4
	3	432±4*	37±2	34±2	25±3	7±2	2.1±0.2	34±3
	4	813±15*	42±3	40±3	33±3	9±2	2.3±0.2	41±4
Red clover	1	51±2	9±1	18±1	6±2	2±1	0,8±0,2	14±2
	2	167±7	20±2	24±2	9±2	4±1	0.9±0.2	28±3
Hungarian sainfoin	1	65±3	15±2	23±2	8±2	4±1	1.0±0.2	13±2
	2	178±6	23±3	35±2	15±2	7±2	1.3±0.2	26±3
	3	269±5*	34±3	39±3	22±3	10±2	1.4±0.2	35±3
	4	311±9*	39±3	47±3	29±3	14±2	1.6±0.2	42±4
White melilot	1	82±4	20±2	18±1	12±2	6±1	3.0±0.2	6±1
	2	260±7*	30±3	31±2	17±2	6±1	3.0±0.2	6±1
Birdsfoot trefoil	1	44±3	18±2	21±2	7±1	3±1	1.1±0.2	33±3
	2	87±6	21±2	25±2	7±1	3±1	1.2±0.2	51±4
	3	101±8	22±2	27±2	11±2	4±1	1.2±0.2	70±4
	4	129±8	24±2	29±2	13±2	5±1	1.4±0.2	89±4
Eastern galega	1	52±4	16±1	28±3	5±2	4±1	1.0±0.2	8±2
	2	108±8	27±2	32±3	11±2	5±1	1.0±0.2	12±2
	3	119±7	31±3	37±3	17±2	9±1	1.1±0.2	16±3
	4	122±8	31±3	42±3	23±3	14±2	1.3±0.2	20±3

*Note: reference data established in accordance with the literature sources listed below

Source: compiled based on the author's own research and [5-7]

An important indicator of the root neck is the number of buds on it, which are able to form a vegetative shoot next year. The largest number of them at the end of the growing season of the first year was observed in the birdsfoot trefoil – 33 pcs., which determines its intensive regrowth and three mowings during the first year of growth. The smallest number of buds was observed on the root neck of white melilot – 6 pcs., which, accordingly, determines its biennial nature and unsatisfactory regrowth with low mowing.

The greatest thickness of the middle part of the central root in the year of sowing was: in white melilot – 6 mm; and the smallest – 2 mm – in red clover. The average thickness of the lateral roots was also the largest in white melilot – 3.0 mm, and the smallest – 0.8 mm – in red clover.

Summing up the findings on the development of root systems of perennial legume grasses in the first year of vegetation, it should be noted: the greatest depth of root penetration into the soil and its greatest

horizontal branching relative to the central root was observed in alfalfa and white melilot. However, the depth of the main mass of root branching in these crops was the smallest; the relatively shallow root system of eastern galega, Hungarian sainfoin, birdsfoot trefoil, and red clover in the year of sowing is compensated by an increase in the depth of the main mass of its branching. In plants of birdsfoot trefoil – also partially by a large radius of horizontal root spread relative to the central root. The greatest thickness of the root neck, central root, and side roots are noted in those types of perennial legumes in which the root system penetrates deeper.

At the end of the second year of vegetation of grasses, the depth of penetration of their roots increased significantly. The root system of white melilot and alfalfa was most deeply penetrated into the soil – up to 260-258 cm. The root system of birdsfoot trefoil was the least deep in the soil – up to 87 cm, which is 3 times less than that of white melilot and alfalfa. There was a similar tendency of root penetration into the soil of

these species of leguminous grasses in depth with the first year of research on grasses that have the largest and smallest depth of penetration into the soil.

The radius of horizontal distribution of root systems of legumes in the second year of vegetation also increased. It was the largest in the plants of white melilot – 30 cm and eastern galega – 27 cm. The lowest horizontal root distribution was observed in red clover and birdsfoot trefoil – 20-21 cm, which is 1.5 times less than in white melilot. Compared to the first year of study, the horizontal distribution of eastern galega roots has significantly increased.

The greatest depth of the main mass of branching of the root system was characteristic of plants of Hungarian sainfoin – 35 cm and eastern galega – 32 cm. The lowest depth of distribution of the main mass of roots was observed in red clover, birdsfoot trefoil and alfalfa – 24-26 cm, which is 1.4 times less than in Hungarian sainfoin.

The thickness of the root neck of perennial legume grasses in the second year of vegetation was 7-18 mm. The thickest root neck was characteristic of alfalfa and white melilot, and the thinnest – for birdsfoot trefoil. The number of buds on the root neck of plants was 6-51 pcs. and it did not depend on the thickness of the root neck. Most of the buds were on the neck of the birdsfoot trefoil, which had the thinnest root neck, and the least – on the root neck of the white melilot, which had the thickest root neck.

The thickness of the central root of legume grasses was 3-7 mm. It was thickest in the Hungarian sainfoin and white melilot, and the thinnest – in the birdsfoot trefoil. To a large extent, the thickness of the central part of the main root was directly proportional to the thickness of the root neck. The thickness of the lateral roots of legume grasses in the second year of vegetation increased slightly and amounted to 0.9–3.0 mm. The thickest were the lateral roots of white melilot plants, and the thinnest were those of red clover.

In the third year of vegetation of perennial legumes, the depth of penetration of their roots into the soil continued to increase and amounted to 100-432 cm. The roots of alfalfa were most deeply penetrated into the soil, and the roots of birdsfoot trefoil were the least deep. The radius of horizontal distribution of the root system of legume grasses was 22-37 cm. It was the largest in alfalfa plants, and the smallest – in birdsfoot trefoil. Compared to previous years of observation, the radius of the root system of alfalfa increases significantly.

The depth of the main mass of branching of the root system of legumes was 27-39 cm at the end of the third year of vegetation. It was largest in the Hungarian sainfoin and eastern galega, and the smallest – in the birdsfoot trefoil. The thickness of the root neck of legume grasses at the end of the third year of vegetation was 11-25 mm. It was the largest in alfalfa plants, and the smallest – in birdsfoot trefoil. The number of buds on the root neck continued to grow and amounted to 16-70 pcs. Most of them were in the birdsfoot trefoil, and the least – in the eastern galega.

The thickness of the central root of legume grasses was 4-10 mm. It was the largest in the plants of Hungarian sainfoin, and the smallest – in birdsfoot trefoil. The thickness of the lateral roots of legume grasses was 1.1–2.1 mm. It was the largest in alfalfa plants, and the smallest – in eastern galega.

In the fourth year of vegetation of perennial grasses, the depth of penetration of their roots was 122-813 cm. The root system of alfalfa was the deepest, and the least deep – in the eastern galega and birdsfoot trefoil. Compared to the previous year, the depth of penetration into the soil of the roots of the birdsfoot trefoil increased and the eastern galega – decreases.

The radius of horizontal distribution of the root system of legume grasses was 22-37 cm. It was the largest in alfalfa plants, and the smallest – in birdsfoot trefoil plants. The depth of the main mass of branching of the root systems of perennial grasses was 29-47 cm. It was the largest in the plants of Hungarian sainfoin, and the smallest – in birdsfoot trefoil.

The thickness of the root neck of plants in the fourth year of vegetation was 13-33 cm. It was the largest in alfalfa plants, and the smallest – in birdsfoot trefoil. The number of renewal buds on the root neck of plants was 20-89 pcs. Most of them were in the birdsfoot trefoil, and the least – in the eastern galega.

The thickness of the central root of legume grasses was 5-14 mm. The thickest root was observed in the plants of Hungarian sainfoin and eastern galega, and the thinnest – in birdsfoot trefoil. Compared to the previous year, the thickness of the central root in eastern galega has significantly increased. The thickness of the lateral roots of plants was 1.3-2.3 mm. It was the thickest in alfalfa plants, and the thinnest – in birdsfoot trefoil and eastern galega.

Dynamics of root system growth and development over all years of study

Analysing the change in the depth of root distribution of perennial legume grasses during all growing years, it was found that during the first two years, the depth of root distribution in all plants was similar. Starting from the third year, there was an intensive growth of alfalfa roots and a slowdown in the root growth of birdsfoot trefoil and eastern galega.

In the second year of vegetation, the depth of penetration of alfalfa roots increased 3.1 times, in the third year – 1.7 times, and in the fourth – 1.9 times. The depth of white melilot roots in the second year increased 3.3 times. The depth of red clover roots in the second year increased by 3.3 times, but their distribution in depth was much less than that of white melilot and alfalfa plants. The roots of Hungarian sainfoin developed similar to the deepening of red clover roots. Their depth in the second year increased by 2.7 times, in the third – 1.5 times, and in the fourth – 1.2 times. The roots of birdsfoot trefoil and eastern galega were least widespread in depth: in the second year, their depth increased by 2.0-2.1 times, in the third year – 1.1-1.2 times, in the fourth – 1.0-1.3 times.

The dynamics of changes in the radius of horizontal distribution of grass roots during all the years of vegetation was more uniform. A significant increase in this indicator was typical for all types of perennial grasses during the first year of vegetation.

A significant increase in the root distribution radius in alfalfa plants occurred in the third year of vegetation – 1.3 times; Hungarian sainfoin plants – in the second and third years of vegetation – 1.5 times each; eastern galega plants – in the second year of vegetation – 1.7 times. A significant increase in the root distribution radius of red clover and white melilot was observed in the second year of vegetation – 1.5 times. The root radius of birdsfoot trefoil plants grew evenly throughout all the years.

The depth of distribution of the main mass of roots in alfalfa plants increased most rapidly: in the second year of vegetation – by 1.7 times, in the third year – by 1.3 times, in the fourth – by 1.2 times. Intensive growth of the depth of the main mass of Hungarian sainfoin roots was observed in the second year of vegetation – 1.5 times and in the fourth year – 1.2 times. The depth of the main mass of white melilot roots in the second year of vegetation increased 1.7 times. The remaining herbs changed the depth of the main mass of roots evenly.

The number of renewal buds on the root neck of perennial legume grasses increased evenly during the growing season. The most intensive increase in their number was characteristic of birdsfoot trefoil: in the second year of vegetation – 1.6 times, in the third year – 1.4 times, in the fourth – 1.3 times. In Hungarian sainfoin and alfalfa plants, the number of buds was similar over the years and increased by 2.1 times in the second year, by 1.4 times in the third year, and by 1.2 times in the fourth year. The number of buds on the root neck of red clover in the second year of vegetation increased by 2 times, and in white melilot, during the two years of vegetation, no changes were observed. The number of buds in eastern galega plants increased 1.3-1.5 times each year.

CONCLUSIONS

Based on the conducted studies and comparisons of biometric indicators of root systems of various types of perennial legumes, the following conclusions can be made:

- the root system of alfalfa is characterised by the largest: penetration into the soil, spread of the roots in a horizontal direction; thickness of the root neck during all the years of vegetation, thickness of the lateral roots during the third or fourth years of vegetation, but the smallest distribution in the soil of the main mass of root branching in the year of sowing and the following

year of vegetation. Such features of the root system development ensure the plasticity, durability, and productivity of alfalfa;

- the root system of white melilot is characterised by the largest: penetration into the soil, thickness of the root neck, central root and lateral roots during all the years of vegetation, horizontal spread of roots in the second year of vegetation, but the smallest number of renewal buds on the root neck during all the years of vegetation. These features determine the high drought and frost resistance of white melilot;

- the root system of the birdsfoot trefoil has the largest number of renewal buds on the root neck during all the years of vegetation. However, the plant had the smallest penetration into the soil, thickness of the root neck and the central root during all the years, radius of horizontal root spread and the depth of the main mass of root branching in the second or fourth years of vegetation, which determines its productive longevity and ability to grow on low-productive and acidic soils;

- the root system of eastern galega is characterised by the largest: depth of the main mass of roots in the soil in all years, the horizontal spread of roots in the second year of vegetation, the thickness of the central root in the fourth year, but the smallest: depth of roots in the soil in the fourth year, thickness of the root neck during the first year and thickness of the side roots during the second, third, and fourth years of vegetation, the number of renewal buds on the root neck during the third and fourth years of vegetation. Such characteristics determine the productive longevity and high biological plasticity of eastern galega;

- the root system of Hungarian sainfoin was characterised by the greatest: horizontal spread of roots and the deepest placement of the main mass of root branching in the third or fourth years of vegetation; thickness of the central root during the second to fourth years. Thus, Hungarian sainfoin plants form a powerful root system in the third and fourth year of vegetation, which determines high biological resistance to adverse growing conditions;

- the root system of red clover has the smallest radius of horizontal root distribution, the thickness of the central root and the thickness of the lateral roots during all the growing years, which affects the possibility of its cultivation in the field crop rotation.

Expanding knowledge about the biological and morphological features of the distribution of root systems of perennial legume grasses during the growing season and in comparison with each other would allow the researchers to understand the processes occurring in plants, as well as select the optimal types of legumes for certain purposes in the context of climate change.

REFERENCES

- [1] Blagoveshchensky, G.V. (2008). Feed production of the Non-Black Soil Zone in a changing climate. *Feed Production*, 10, 6-8.
- [2] Gaitov, T.A. (2001). Perennial grasses as a factor in improving the efficiency of land use. *Feed Production*, 8, 16-18.
- [3] Korniyuchuk, O.V., Buhayov, V.D., Hetman, N.Ya., Satanovska, I.P., Tsyhansky, V.I., Kvitko, H.P., & Protopish, I.H. (2013). The role of alfalfa sowing in the intensification of feed production. *Handbook of Ukrainian Farmers*, 2, 222-225.

- [4] Brun, I.M. (2007). Influence of weather factors on the growth, development and formation of the crop of the leaf stem mass of the espresso of sand in the conditions of the right bank of the Forest-steppe. *Feed and Feed Production*, 59, 71-76.
- [5] Kvitko, H.P. (2010). Adaptive energy-saving technologies for growing perennial legumes for fodder in the Forest-Steppe of the Right Bank. *Feed and Feed Production*, 66, 78-82.
- [6] Makarenko, P.S. (2008). *Meadow and field fodder production*. Vinnytsia: Private individual Danyliuk VG.
- [7] Kirilesko, O.L. (2013). Influence of saturation of fodder crop rotations with perennial grasses and intermediate crops on the balance of humus in the soil. *Feed and Feed Production*, 76, 151-157.
- [8] Zabarna, T.A. (2009). Formation of leaf-stem and root mass of meadow clover in the second year of life in the conditions of the right-bank Forest-Steppe of Ukraine. *Feed and Feed Production*, 64, 148-155.
- [9] Schnidtko, K., & Rauber, R. (1990). Gefardet der Leduminosenanbau im ökologischen Landbau die Grundwassergualitat? *Bio-Land*, 5, 15-18.
- [10] Ghosh, P.K., & Maiti, T.K. (2016). Structure of extracellular exopolysaccharides (EPS) produced by rhizobia and their functions in legume-bacteria symbiosis: A review. *Achievements in the Life Sciences*, 10, 136-143.
- [11] Marczak, M., Mazur, A., Koper, P., Zebracki, K., & Skorupska A. (2017). Synthesis of rhizobial exopolysaccharides and their importance for symbiosis with legume plants. *Genes*, 8(12), 360.
- [12] Stevens, G.G., Perez-Fernandez, M.A., Morcillo, R.J.L., Kleinert, A., Hills, P., Brand, D.J., Steenkamp, E.T., & Valentine, A.J. (2019). Roots and nodules response differently to P starvation in the mediterranean-type Legume *Virgilia divaricata*. *Frontiers in Plant Science*, 10, article number 73. doi: 10.3389/fpls.2019.00073.
- [13] Suzaki, T., Takeda, N., Nishida, H., Hoshino, M., Ito, M., Misawa, F., Handa, Y., Miura, K., & Kawaguchi, M. (2019). Lack of symbiont accommodation controls intracellular symbiont accommodation in root nodule and arbuscular mycorrhizal symbiosis in *Lotus japonicus*. *PLOS Genetics*, 15(1), article number e1007865. doi: 10.1371/journal.pgen.1007865.
- [14] Jacob, C., Carrasco, B., & Schwember, A.R. (2016). Advances in breeding and biotechnology of legume crops. *Plant Cell Tissue Organ Culture*, 127(3), 561-584. doi: 10.1007/s11240-016-1106-2.
- [15] Adams, M.A., Buchmann, N., Sprent, J.I., Buckley, T.N., & Turnbull, T.L. (2018). Crops, nitrogen, water: Are legumes friend, foe, or misunderstood ally? *Trends in Plant Science*, 23(6), 539-550. doi: 10.1016/j.tplants.2018.02.009.
- [16] Kvitko, G.P., Tkachuk, O.P., & Hetman, N.Ya. (2012). Perennial legumes are the basis of natural intensification of fodder production and improvement of soil fertility in the Forest-Steppe of Ukraine. *Feed and Feed Production*, 73, 113-117.
- [17] Moiseychenko, V.F., & Yeshchenko, V.O. (1994). *Fundamentals of scientific research in agronomy*. Kyiv. Vyshcha shkola.
- [18] Babych, A.O. (1994). *Methods of conducting experiments on feed production*. Vinnytsia.

БІОЛОГІЧНІ ОСОБЛИВОСТІ ПОШИРЕННЯ КОРЕНЕВИХ СИСТЕМ БОБОВИХ БАГАТОРІЧНИХ ТРАВ В УМОВАХ ЗМІНИ КЛІМАТУ

Олександр Петрович Ткачук

Вінницький національний аграрний університет
21008, вул. Сонячна, 3, м. Вінниця, Україна

Анотація. Визначальним чинником при виборі бобових багаторічних трав за умов змін клімату є їх посухостійкість, адже розвиток коренів прямо впливає на цю властивість, а також визначає їх продуктивність. Метою досліджень було встановити, за рахунок яких морфологічно-біологічних характеристик коренів трав забезпечуються господарські особливості в умовах змін клімату. Застосовували методи спостереження, порівняння, польового досліду. З'ясовано, що корені люцерни посівної мають найбільше заглиблення у ґрунт, поширення у горизонтальному напрямі, товщину кореневої шийки та бічних корінців, що забезпечує пластичність, довговічність і продуктивність її посівів. Встановлено, що корені буркуну білого мають найбільше заглиблення у ґрунт, товщину кореневої шийки, центрального кореня і бічних корінців, їх поширення у горизонтальному напрямі, що впливає на посухо- і морозостійкість рослин. Обґрунтовано, що корені лядвенцю рогатого відзначаються найбільшою кількістю бруньок відновлення на кореневій шийці, найменшим заглибленням, товщиною кореневої шийки, що впливає на продуктивне довголіття та можливість росту на бідних і кислих ґрунтах. Виявлено, що корені козлятника східного мали найбільше заглиблення основної маси, поширення у горизонтальному напрямі, товщину центрального кореня, що впливає на продуктивне довголіття і високу біологічну пластичність. Встановлено, що корені еспарцету піщаного відзначались найбільшим поширенням у горизонтальному напрямі та найглибшим розміщенням основної маси розгалужень, що впливає на високу біологічну стійкість до несприятливих умов вирощування. З'ясовано, що корені конюшини лучної мали найменший радіус поширення у горизонтальній площині, товщину центрального і бічних корінців, що визначає можливість її вирощування у польовій сівозміні. Зазначені наукові знання дозволять підібрати види бобових багаторічних трав для повної реалізації їх потенціалу за умов зміни клімату

Ключові слова: рослини, корені, підземний розвиток, морфологія, роки вегетації

**Журнал
«НАУКОВІ ГОРИЗОНТИ»**

**Том 24, № 2
2021**

(Англійською мовою)

Редагування англomовних текстів:

С. Воровський, А. Кравченко

Літературний редактор:

С. Пастух

Редагування бібліографічних списків:

С. Пастух, К. Сосєдко

Комп'ютерна верстка:

К. Сосєдко

Підписано до друку з оригінал-макета 23.06.2021
Ум. друк. арк. 14,6

Видавництво Поліський національний університет
10008, б-р Старий, 7, м. Житомир, Україна.
Тел. (0412) 22-04-17
E-mail: info@sciencehorizon.com.ua
www: <https://sciencehorizon.com.ua>

**Journal
"SCIENTIFIC HORIZONS"**

**Volume 24, No. 2
2021**

Editing English-language texts:

S. Vorovsky, A. Kravchenko

Literary editor:

S. Pastukh

Editing bibliographic lists:

S. Pastukh, K. Sosiedko

Desktop publishing:

K. Sosiedko

Signed to the print with the original layout 23.06.2021.

Mind. print. ark. 14.6

Publisher: Polissia National University
10008, 7 Staryi Blvd., Zhytomyr, Polissia National University, Ukraine.
Tel. (0412) 22-04-17;
E-mail: info@sciencehorizon.com.ua
www: <https://sciencehorizon.com.ua/en>