

MODERN PHYTOMORPHOLOGY



ISSN 2226-3063
e-ISSN 2227-9555



Editorial Team
Executive Editor

Andrew Novikoff (Andriy Novikov) State Natural History Museum NASU, Lviv, Ukraine

Editorial Assistant Sergii Kondratyuk M.G. Kholodny Institute of Botany NASU, Kyiv, Ukraine

Editorial Board Ali Raza Fujian Agriculture and Forestry University (FAFU), Fuzhou 350002, China

SUN Shucun Professor of Ecology, Department of Biology Nanjing University, China

Mehmet Karaca Akdeniz University, Turkey

Yosyp Berko S.Z. Gzhytskyj Lviv National University of Veterinary Medicine and Biotechnologies, Lviv, Ukraine

Vasyl Budzhak Yuriy Fedkovich Chernivtsi National University, Chernivtsi, Ukraine

Lyudmila Bukhtiyarova Institute for Evolutionary Ecology of the NAS of Ukraine Kyiv, Ukraine

Roland Eberwein Carinthian Botanic Center Klagenfurt am Woerthersee, Austria

Olexandr Klymyshyn State Natural History Museum NASU, Lviv, Ukraine

Vladislav Korzhenevsky Nikitsky Botanical Gardens, National Scientific Centre Yalta, Ukraine

Joanna Korzeniak Institute for Nature Conservation PAS Cracow, Poland

Zvenyslava Mamchur Ivan Franko Lviv National University, Lviv, Ukraine

Anastasia Odintsova Ivan Franko Lviv National University, Lviv, Ukraine

Olga Terek Ivan Franko National University of Lviv, Lviv, Ukraine

Antonio Tiezzi Tuscia University, Viterbo, Italy

R. Sakthivel, Ph.D. Principal St. Eugene University, Zambia, Zambia

Yurii Chernobay State Natural History Museum NASU, Lviv, Ukraine

Illia Chornej Yuriy Fedkovich Chernivtsi National University, Chernivtsi, Ukraine

Alexey Shipunov Minot State University, Minot, USA

Svitlana Shevchenko Nikitsky Botanical Garden
National Scientific Centre, Yalta, Ukraine

Kazimierz Szczepanek Institute of Botany PAS, Cracow, Poland

AYSE GUL INCE Vocational School of Technical Sciences
Akdeniz University, Turkey

Articles in press and Articles in process

Opinion

[The Crucial Connection: Soil Drainage's Impact on Plant Growth](#)

Kaneeka Banyal, Pooja Sharma

107-108

Research Article

[Determining the effectiveness of ozone in prolonging the preservation of soft cheese](#)

Safa Abdullah Azzawi*, Ziad Tariq Samir and Mohammed Ahmed Jassim

89-94

Research

[Effect of cutting irrigation at different stages of growth and spraying with bitter melon extract on some vegetative qualities of two varieties of Zea mays L.](#)

Tamara Sabah Hadi, Wisam Malik Dawood, Mohammed Ali Abood

89-97

Research Article

[Modification of Seeds to Roots: The Uniqueness of Germination and Growth of Putat \(*Barringtonia acutangula*\)](#)

Pinta Murni, Bambang Hariyadi, Mahya Ihsan, Agus Subagyo, Winda Dwi Kartika

80-86

Research

[The efficiency of urea-ammonium nitrate application in inter-row feeding in maize cultivation](#)

Adamchyk Y., Kravchenko N., **Kolisnyk O.**, **Aralova. T.**, Protasov O., Dubovyk O., Dubovyk I., Stavytskyi A.

113-117

Research Article

[Comparative flower morphology in *Allium sativum* L. \(Amaryllidaceae\).](#)

O. Fishchuk, N. Kostruba, N. Chmil

109-112

Research

[Criteria for evaluating the state of rare plant species populations](#)

Klymenko H., Artemenko D., Klymenko I., Kovalenko N., Melnyk A., Butenko S., Horbas S., Tovstukha O.

98-106

Research Article

[Using regional normalized difference vegetation index for the large-scale yield prediction of potato, vegetables, fruits, and berries, cultivated in Kherson region of Ukraine](#)

Pavlo Lykhovyd, Raisa Vozhehova, Oleksandr Averchev, Oleksandr Rudik, Liudmyla Hranovska, Sergiy Lavrenko, Nataliia Avercheva, Grigoriy Latiuk

118-124

Research Article

[Technological methods of improving rapeseed feed and reducing their toxicity](#)

Mykytyn Mykola, Melnyk Uliana, Hotvianska Anna, Kovalenko Vitalii, Bondarenko Oksana, Bordun Roman

125-133

Brief Report

[Study on Phenology of Some Papilionoid taxa \(Papilionoideae- Leguminosae\).](#)

RACHITHA

72-75

Research Article

[Annual NDVI dynamics observed in industrial tomato grown in the south of Ukraine](#)

Pavlo Lykhovyd*, Raisa Vozhehova, Sergiy Lavrenko

DOI: 10.5281/zenodo.10012129

164-169

Research Article

Popularization of Finger Millet Production, Value Addition, and Marketing in
Wayu Tuka Were da, East Wollega Zone

Amanuel Berhanu Bonso

DOI: 10.5281/zenodo.10012175

83-92

Short Communication

A preliminary study on the effects of pre-planting heating o saffron corms
germination speed and number of sprouts in greenhouse conditions

Pavlo Lykhovyd, Olesia Yuziuk, Oleksandr Rudik, Olena Ozhovan, Liudmyla
Popova, Natalia Rudik, Valerii Dubrovin

179-181

Review Article

Overview of Obesity and Pharmacological Significance of Sita Ashoka (Saraca
asoca) Helpful in management of obesity and other Diseases: A Review

Shivam Vashist, Kamal, Pooja Rathi, Anish, Rohit, Dhirender Kaushik, Shivkant
Sharma

201-212

Research Article

The genus *Thrixspermum* Loureiro in Arunachal Pradesh and three new additions

Khyanjeet Gogoi, K. Chowlu

182-186

Case Report

A Case Study on the Use of Hydrogen Peroxide as an Agent for Radish Seeds
Germination Promotion

Pavlo Lykhovyd, Raisa Vozhehova , Mykhailo Fedorchuk, Valentyna Fedorchuk
Mykhailo Kobelev

213-216

[Submit Manuscript](#)

Article Type: Research
J Name: Modern Phytomorphology
Short name: MP
ISSN: ISSN 2226-3063/ eISSN 2227-9555
Year: 2023
Volume: 17
Page numbers: 113-117
DOI: 10.5281/zenodo.200121
(10.5281/zenodo.Year-Volume-PDFNo.)
Short Title: The efficiency of urea-ammonium nitrate application in inter-row feeding in maize cultivation

RESEARCH ARTICLE

The efficiency of urea-ammonium nitrate application in inter-row feeding in maize cultivation

Adamchyk Y.^{1*}, Kravchenko N.¹, Kolisnyk O.², Aralova. T.², Protasov O.¹, Dubovyk O.¹, Dubovyk I.¹, Stavtyskiy A.³

¹Sumy National Agrarian University, H. Kondratieva St., 160, Sumy, 40021, Ukraine.

²Vinnitsia National Agrarian University, Soniachna Str. 3, 21008, Vinnitsia, Ukraine

³Separate Structural Subdivision, Okhtyrka Professional College of Sumy National Agrarian University, Sumy region, Okhtyrka, Sumska Str., 46, 42700, Ukraine

*Corresponding author: Adamchyk Y., Sumy National Agrarian University, Gerasim Kondratyev Str. 160, 40021, Sumy, Ukraine; Email: andb201727@ukr.net

Received: 06.10.2023, Manuscript No.: mp-23-115023 | Editor Assigned: 13.10.2023, Pre-QC No. mp-23-115023(PQ) | Reviewed: 19.10.2023, QC No. mp-23-115023(Q) | Revised: 26.10.2023, Manuscript No. mp-23-115023(R) | Accepted: 01.11.2023 | Published: 09.11.2023

Abstract

The issue of influence on the level of grain maize yield of different rates of application of Urea Ammonium Nitrate (UAN-32) in the BBCH 17-18 phase is highlighted. The fertilizer was applied by inter-row cultivation with a mounted cultivator IRIS-T. The most optimal rate of nitrogen fertilizer application during root application is N_{35} , in conditions of dark gray soil, on loess loam in the Polissya zone of Ukraine, against the background of applied nitrogen N_{98} in autumn and when sowing N_8 as part of a complex fertilizer (YaraMila 8-24-24). The application rate of UAN above N_{35} reduces grain yield and profitability. Applied during the growing season, the norms of N_{16} - N_{35} of nitrogen fertilizer contribute to an increase in the height, weight of the plant, leaf surface area and grain yield.

Keywords: fertilizer rates, nitrogen, yield, productivity, corn, inter-row cultivation, UAN (Urea Ammonium Nitrate).

Introduction

Nitrogen application is one of the key points in the grain maize fertilization system. Numerous works of researchers give examples of the use of nitrogen fertilizers in different forms, at different times, and methods of application (Kharchenko et al., 2021a; Simon et al., 2022). But there are still discussions about the rapid availability of ammonium and nitrate nitrogen to the plant (Lopushniak et al., 2021). A number of scientists talk about the significant impact on the development of maize is the application of nitrogen fertilizer in pre-sowing tillage (Zakharchenko et al., 2023a). There is also information about the positive effect of nitrogen application in the composition of pre-sowing complex fertilizer for the optimal beginning of the growing season and the biological cycle. But in the fertilization system, the importance of foliar feeding and, in recent times, also root nutrition is distinguished. It is known that the maize plant has critical phases of development and at this time the signs of physiological starvation for certain nutrients, in particular, nitrogen, are too acute (Zakharchenko et al., 2023b; Zhang et al., 2023). Therefore, nitrogen application during the growing season is carried out mainly in the BBCH 13-17 phases. The nitrogen supply of the plant also depends on the type of soil (Zanella et al., 2022), place in the crop rotation (Mishchenko et al., 2019; Ivanov et al., 2019;

Palamarchuk et al., 2021a), weather and climatic conditions (Kharchenko et al., 2021b) the location of the field/plant on the ground (Dindaroglu et al., 2021), and the tillage system.

The application of nitrogen fertilizers affects the increase in plant height, leaf surface area, chlorophyll content, plant weight, grain weight per cob and, accordingly, a higher yield. The quality of the grain also depends on the application of nitrogen fertilizers.

The desire for super-high grain yields leads to the use of high doses of nitrogen, especially during fertigation, but this can cause nitrogen losses in the form of nitrates or nitrogen gases. The addition of urease inhibitors significantly reduces ammonia evaporation. Application of UAN, together with nitrification inhibitors and urease, increases the available nitrogen content for the plant and reduces the loss of nitrate nitrogen. Both UAN alone and in combination with the above-mentioned compounds, there is an increase in the nitrogen content in the grain, in general the grain yield (Palamarchuk et al., 2018; Ren et al., 2023). But there are other opinions that it is better not to mix UAN with inhibitors (Li et al., 2023).

Materials and Methods

The research was carried out in production conditions on the territory of the farm of LLC "State Enterprise "Zernyatko" of the Koryukiv district of the Chernihiv region. The geographical coordinates of the location of the plots are 51.549364,32.375249. The field is dominated by dark gray podzolic medium-loamy soil on loess loam. Total humus content is 3.05 % (according to Tyurin), pHKCl 5.04, pHH₂O 5.76, NO₃+NH₄+12.6 mg kg⁻¹ (very low), mobile phosphorus 77.8 mg kg⁻¹ according to Machygin (medium level), exchangeable potassium 49.5 mg/kg according to Kirsanov (medium level), sulfur content 15.95 mg kg⁻¹ (very high), exchangeable calcium 5.99 mg-eq/100 g according to Schollenberg (medium), magnesium 1.01 mEq/100 g according to Schollenberg (medium). The content of microelements is low: zinc 1.32 mg kg⁻¹, copper 0.13 mg kg⁻¹, manganese 6.38 mg kg⁻¹ (atomic absorption spectrometry method), boron 0.19 mg kg⁻¹ (hot water determination). Since the soil pH is slightly acidic, there is a decrease in the level of absorption of phosphorus, potassium, nitrogen and these microelements. Liming has not been carried out for twenty years. The area of the accounting plot is 50 m² with the three replications. The research was conducted in 2021-2022. The predecessor of maize in field crop rotation is sunflower. UAN-32 was used in inter-row fertilization in the development phase of BBCH 17-18 maize. Inter-row fertilization was carried out by the IRIS-T mounted unit in combination with the John Deere 6135 tractor. The Pioneer 8816 maize hybrid was used in the experiment, the results of the experiment were recorded after threshing each plot and weighing threshed crop on truck scales. Moisture during harvesting was 22%-23% and was converted to 14% moisture when calculating the crop.

The scheme of the experiment is as follows:

1. Control (without inter-row fertilization UAN).
2. Inter-row fertilization UAN-32 50 l ha⁻¹ N16.
3. Inter-row fertilization UAN-32 80 l ha⁻¹ N25.6.
4. Inter-row fertilization UAN-32 110 l ha⁻¹ with N35.
5. Inter-row fertilization UAN-32 140 l ha⁻¹ N44.8.

Biometric indicators, namely the measurement of plant height and leaf surface area, were carried out in the phase of milky ripeness of the plant. The leaf surface area was determined by measuring the length and width of the leaf and multiplying by a conversion factor of 0.67. The results of the analysis were calculated statistically using Agrostat program.

The main cultivation of the field was carried out in a complex with the application of nitrogen fertilizer with a unit for the application of anhydrous ammonia 120 kg ha⁻¹ to a depth of 20 cm-22 cm. Sowing of maize was carried out on April 20-25, 2021-2022 with a norm of 75 000 plants ha⁻¹ to a depth of 3 cm-4 cm. Sowing was carried out with a Kinze 3600 tilled sixteen-row seeder, with a row spacing of 70 cm. Along with sowing, Yara Mila 8-24-24 complex fertilizer was applied at a rate of 100 kg ha⁻¹ in physical weight.

Results and Discussion

Tab. 1 shows the results of measuring the height of the plant and the leaf surface area in the phase of milky-wax ripeness. As you can see, the plants react to the applied fertilizer and with an increase in the nitrogen rate in the UAN, the

height of the plant increases. This is a proven fact of the effect of nitrogen on plants. There is a significant difference between the variants and a direct correlation with the increase in the norm. The results of determining the leaf surface area of the maize plant depending on the applied doses of UAN in inter-row cultivation also confirm our previous statements and the stable conclusions of agrochemists that nitrogen fertilizers have a significant impact on the above-ground mass, on the development of photosynthetic potential that can maximize the use of solar energy.

Table 1. Biometric indicators of maize plants (average for 2021-2022)

Variant	Plant height, cm	Leaf surface area, thousand m ² ha ⁻¹
1. Control (without applying UAN to inter-row cultivation)	246.7	46.8
2. UAN-N16	250.2	47.2
3. UAN-N25.6	252.4	47.4
4. UAN-N35	254.1	47.5
5. UAN-N48	256.5	47.8
LSD ₀₅	0.9	0.3

Since the application of UAN was carried out in the root zone in the phase of 7-8 leaves, there was no difference in the dates of the onset of the phase of 11 leaves, panicle ejection, milky and wax ripeness and is not given in the article.

Scientists of Vinnytsia National Agrarian University also emphasize that foliar and root fertilization of maize affect not only the height of the plant and the area of the leaf apparatus, but also the height of the cob attachment, the length of the cob leg, and the number of cob wrappers (Palamarchuk & Solomon, 2021b). Scientists also note that the effect of fertilizer has a different effect on maize hybrids with different FAO, in them it was the early-ripening hybrid that had the highest response to the applied fertilizers – Ecolist, Rostok, Vympel.

In some cases, the biometric indicators may be lower than at the control. For example, if the research is aimed at studying the effects of soil pollutants on plant growth, development and grain yield. Scientists of the National Scientific Center "Institute of Agriculture of the National Academy of Agrarian Sciences of Ukraine" (Korsun et al., 2016) note that when polluted, all parameters of the photosynthetic potential of the plant decrease by 5, 10 and 100 times.

Tab. 2 shows the weight of the leaf-stem mass of one maize plant and the weight of grain from the cob as one of the important elements of nitrogen fertilizer application efficiency. It is the ratio of the main and by-products that reflects the effect of the applied fertilizer and other factors influencing the yield of the crop. (Tab. 2).

Table 2. Structural elements of maize plant yield

Experiment Option	Weight of leaf-stem mass and cobs of 1 plant, kg	Grain weight per cob, kg	Yield, t ha ⁻¹
1. Control (without applying UAN to inter-row cultivation)	0.435	0.131	12.82
2. UAN-N16	0.491	0.142	13.21
3. UAN-N25.6	0.515	0.152	13.42
4. UAN-N35	0.529	0.164	13.66
5. UAN-N48	0.545	0.162	13.60
LSD ₀₅	0.013	0.012	0.25

The weight of the leaf-stem mass together with the cobs is 0.435 kg-0.545 kg and is directly proportional to the applied UAN rate. We cannot say the same about the mass of grain on the cob. There is a significant difference between the variants, with the exception of a UAN of 110 kg ha⁻¹ and a UAN of 140 kg ha⁻¹, between which there is no difference. That is, with the acquisition of a greater weight of the leaf-stem mass, the weight of the grain from the cob decreased.

According to the study, we observe, as expected, that in the control area, the yield of grain maize is the lowest among other studied variants and amounts to 12.82 tha⁻¹. The reason is that no additional nitrogen rates were received in this area. The next plot with a root fertilization rate of 50 kg ha⁻¹ of the physical weight of UAN

fertilizer after harvesting and its calculation showed a yield of 13.21 t ha⁻¹, which is 0.39 tons more than in the control. The experimental plot with a fertilization rate of 80 kg ha⁻¹ gave a yield of 13.42 t ha⁻¹, which is relatively higher than the previous one and gave an increase in yield relative to the control by 0.6 t ha⁻¹. The largest relative increase was given by the plot with the UAN application rate of 110 kg ha⁻¹—0.84 t. The experiment with the norm of 140 kg ha⁻¹ of nitrogen fertilizer, although it showed an increase in yield, which was 0.78 t, did not continue the pattern of the experiment to increase the yield with an increase in the fertilizer rate. Thus, the study showed that an increase in the application rates of UAN in the root fertilization of maize is accompanied by an increase in yield, but stops increasing it after an increase in the application rate of more than 110 kg ha⁻¹. These results can be explained by physiological processes in plants, as well as by the rules of agriculture. Plants always absorb only those elements that are necessary for their life, and they do so only in the quantities they need. They are not able to replace some elements with others, since each plays its own role in biochemical processes. Turning to the rules of agriculture established by Justus Liebig, we find an explanation for the results of the experiment, namely, the law of optimum, which explains that the greatest yield can be obtained under conditions of optimal concentrations of elements available to plants. Because the closer the optimal concentration of all the necessary components for plants is, the more efficiently they are used, which is reflected in the increase in yield. Therefore, the rate of 110 kg ha⁻¹ of UAN, which in terms of the active ingredient is 35 kg ha⁻¹ of nitrogen, creates the most optimal concentration of elements necessary for maize plants. An increase in the rate to 140 kg ha⁻¹, equivalent to 45 kg ha⁻¹ of nitrogen for the active substance, led to a further decrease in yield, as the concentration shifted again.

In this experiment, it is also appropriate to talk about the principles of antagonism and synergy of ions. Antagonism can be explained by the decrease in yield gain after application of a rate greater than 110 kg ha⁻¹, since the excess of nitrogen compounds led to competition of elements when absorbed by the roots and led to a greater deficiency of others in plants. Also, at the norm that made it possible to obtain the highest yield, the phenomenon of synergy manifested itself, when, at optimal norms of the content of elements, they increased the effect of each other.

Such data confirm the work of a number of scientists. For example, in North Carolina, when UAN was applied at different doses from 39 kg ha⁻¹ to 235 kg ha⁻¹, the yield increased significantly (Cahill et al., 2007). Chinese scientists investigated the effectiveness of adding phosphorus, magnesium and sulphur to UAN in the cultivation of grain maize in different doses and rates. The main factor influencing the effectiveness of a particular fertilizer was weather conditions. A good effect was obtained with pre-sowing application of urea mixture with phosphorus and then in top feeding together with sulfur, the increase in grain yield was 29%-35% (Li et al., 2021). Other scientists have also noted that the use of nitrogen in the cultivation of maize above 300 kg ha⁻¹ is excessive and can lead to a decrease in the utilization rate of nitrogen and other elements and can cause environmental problems such as eutrophication of water bodies and global warming (Zhang et al., 2015; Good & Beatty, 2011).

The use of nitrogen fertilizers, in particular, urea ammonium nitrate, increases the yield of grain maize, but in a specific natural and climatic zone, it is necessary to select the appropriate scientifically based dose and timing of fertilizer application for the best effect and profitable economic efficiency.

Conclusions

The application of UAN has a positive effect on the height of maize plants and the area of the leaf surface. The level of exposure depends on the dose of nitrogen application. With an increase in the rate of nitrogen, the weight of the leaf-stem mass and grain increases. The introduction of urea ammonium nitrate with inter-row tillage in root fertilization in the phase of 7-8 leaves increases the yield of grain maize. The increase in yield increases approximately uniformly with an increase in the application rate of UAN, therefore, with limited resources, its application is also advisable, since it also brings additional income. In the conditions of dark gray forest soils of the Chernihiv Polyssia of Ukraine, it is recommended to use UAN-32 on maize crops with inter-row cultivation in the root application with rates of up to 110 kg ha⁻¹ in the phase of 7-8 leaves to increase yield.

References

- Cahill S., Osmond D., Crozier C., Israel D., Weisz R. 2007. Winter Wheat and Maize Response to Urea Ammonium Nitrate and a New Urea Formaldehyde Polymer Fertilizer. *Agron. J.*, **99**: 1645-1653.
- Dindaroglu T., Tunguz V., Babur E., Alkharabsheh H.M., Seleiman M.F., Roy R., Zakharchenko E. 2022. The use of remote sensing to characterise geomorphometry and soil properties at watershed scale. *Int. J. Global Warming*, **27**: 402-421.
- Good A.G., Beatty P.H. 2011. Fertilizing Nature: A Tragedy of Excess in the Commons. *PLoS Biol.*, **9**: e1001124.
- Hryhoriv Ya.Ya., Butenko A.O., Moisiienko V.V., Panchyshyn V.Z., Stotska S.V., Shuvar I.A., Kriuchko L.V., Zakharchenko E.A., Novikova A.N. 2021. Photosynthetic activity of *Camelina sativa* plants depending on technological measures of growing under conditions of Precarpathians of Ukraine. *Mod. Phytomorphology*, **15**: 17-21.
- Ivanov M.I., Rutkevich V.S., Kolisnyk O.M., Lisovyi I.O. 2019. Research on the block-portion separator parameters influence on the adjustment range of operating elements speed. *INMATEH Agric. Eng.*, **57**: 37-44.
- Kharchenko O., Petrenko S., Sobko M., Medvid S., Zakharchenko E., Pschyhenko O. 2021b. Models of quantitative estimation of sowing density effect on maize yield and its dependence on weather conditions. *Sci Papers Ser A Agron.*, **2**: 224-231.
- Kharchenko O.V., Petrenko S.V., Sobko M.G., Medvid S.I., Zakharchenko E.A. 2021a. Nutrients use efficiency by modern hybrids of maize under arid conditions of the Forest-Steppe. *Agrochem. Soil Sci.*, **91**: 49-58.
- Korsun S., Buslayeva N., Dovbash N. 2016. Features of photosynthetic activity of sowings of corn for grain in conditions of pollution with lead, cadmium, and zinc. *Herald of Agrar. Sci.*, **1**: 32-36.
- Li X., Zhang X., Wang S., Hou W., Yan L. 2023. The Combined Use of Liquid Fertilizer and Urease/Nitrification Inhibitors on Maize Yield, Nitrogen Loss and Utilization in the Mollisol Region. *Plants*, **12**:1486.
- Li H., Mei X., Nangia V., Guo R., Liu Y., Hao W., Wang J. 2021. Effects of different nitrogen fertilizers on the yield, water- and nitrogen-use efficiencies of drip-fertigated wheat and maize in the North China Plain. *Agric. Water Manage.*, **243**: 106474.
- Lopushniak V., Hrytsuliak H., Gamayunova V., Kozan N., Zakharchenko E., Voloshin Y., Lopushniak H., Polutrenko M., Kotsyubynska Y. 2022. A Dynamics of macro elements content in eutric podzoluvisols for separation of wastewater under Jerusalem artichokes. *J. of Ecol. Eng.*, **23**: 33-42.
- Mishchenko Y.G., Zakharchenko E.A., Berdin S.I., Kharchenko O.V., Ermantraut E.R., Masyk I.M., Tokman V.S., Kharchenko O.V. 2019. Herbolological monitoring of efficiency of tillage practice and green manure in potato agroecosystem. *Ukrainian J. of Ecol.*, **9**: 210-219.
- Palamarchuk V., Honcharuk I., Telekalo N., Krychkovskiy V., Kupchuk I., Mordvaniuk M. 2021a. Modeling of hybrid cultivation technology corn to ensure energy efficiency for sustainable rural development. *Ukrainian J. of Ecol.*, **11**: 204-211.
- Palamarchuk Vitalii, Honcharuk Inna, Honcharuk Tetiana, Telekalo Natalia. 2018. Effect of the elements of corn cultivation technology on bioethanol production under conditions of the right-bank forest-steppe of Ukraine. *Ukrainian J. of Ecol.*, **8**: 47-53.
- Palamarchuk V., Solomon A. 2021b. Research of the corn assimilation surface formation depending on foliar feeding. *Feeds and Feed Prod.*, **92**: 82-94.
- Ren B., Huang Z., Liu P., Zhao B., Zhang J. 2023. Urea ammonium nitrate solution combined with urease and nitrification inhibitors jointly mitigate NH₃ and N₂O emissions and improves nitrogen efficiency of summer maize under fertigation. *Field Crops Res.*, **296**: 108909.
- Simon A., Russu F., Ceclan A., Popa A., Bardas M., Chetan F., Rusu T. 2022. Mineral fertilization- an important factor in obtaining maize harvests. *Agrorlife Sci. J.*, **11**: 218-225.
- Zakharchenko E., Datsko O., Mishchenko Y., Melnyk A., Kriuchko L., Rieznik S., Hotvianska A. 2023a. Efficiency of biofertilizers when growing corn for grain. *Mod. Phytomorphology*, **17**: 50-56.
- Zakharchenko E.A., Petrenko S.V., Berdin S.I., Podhaietskyi A.A. Kravchenko N.V., Hnitetskyi M.O., Hlupak Z.I., Bordun R.M., Tiutiunnyk O.S., Tryus V O. 2023b. Response of maize plants to seeding rates under conditions of typical black soil. *Mod. Phytomorphology*, **17**: 71-74.
- Zanella A., Ponge J.-F., Jabiol B., Van Delft B., De Waal R., Katzensteiner K., Kolb E., Bernier N., Mei G., Blouin M., Juilleret J., Pousse N., Stanchi S. Cesario, Le Bayon R.-C., Tatti D., Chersich S., Carollo L., Englisch M., Schrötter A., Schauffer J., Bonifacio E., Fritz I., Sofo A., Bazot S., Lata J.-C., Iffly J.-F., Wetzel C.E., Hissler C., Fabiani G., Aubert M., Vacca A., Serra G., Menta C., Visentin F., Cools N., Bolzonella C., Frizzera L., Zampedri R., Tomasi M., Galvan P., Charzynski P., Zakharchenko E., Waez-Mousavi S.M., Brun J.-J., Menardi R., Fontanella F., Zaminato N., Carollo S., Brandolese A., Bertelle M., Zanella G., Bronner T., Graefe U., Hager H. 2022. A Standardized Morpho-Functional Classification of the Planet's Humipedons. *Soil Syst.*, **6**: 59.
- Zhang X., Wang Q., Xu J., Gilliam F.S., Tremblay N., Li C. 2015. In Situ Nitrogen Mineralization, Nitrification, and Ammonia Volatilization in Maize Field Fertilized with Urea in Huanghuaihai Region of Northern China. *PLoS ONE*, **10**: e0115649.
- Zhang X., Wu D., Zakharchenko E.A. 2022 Review on effects of biogas slurry application on crop growth. *Agrar. Innov.* **13**: 155-166.